

## Review

# Elevated CO<sub>2</sub> and plant structure: a review

SETH G. PRITCHARD,\* HUGO H. ROGERS,\* STEPHEN A. PRIOR\* and CURT M. PETERSON<sup>1</sup>

\*USDA-ARS National Soil Dynamics Laboratory, 411 South Donahue Drive, Auburn, AL 36832, USA, <sup>1</sup>Department of Biological Sciences, University of Northern Colorado, Greeley, CO 80639, USA.

### Abstract

Consequences of increasing atmospheric CO<sub>2</sub> concentration on plant structure, an important determinant of physiological and competitive success, have not received sufficient attention in the literature. Understanding how increasing carbon input will influence plant developmental processes, and resultant form, will help bridge the gap between physiological response and ecosystem level phenomena. Growth in elevated CO<sub>2</sub> alters plant structure through its effects on both primary and secondary meristems of shoots and roots. Although not well established, a review of the literature suggests that cell division, cell expansion, and cell patterning may be affected, driven mainly by increased substrate (sucrose) availability and perhaps also by differential expression of genes involved in cell cycling (e.g. cyclins) or cell expansion (e.g. xyloglucan endotransglycosylase). Few studies, however, have attempted to elucidate the mechanistic basis for increased growth at the cellular level.

Regardless of specific mechanisms involved, plant leaf size and anatomy are often altered by growth in elevated CO<sub>2</sub>, but the magnitude of these changes, which often decreases as leaves mature, hinges upon plant genetic plasticity, nutrient availability, temperature, and phenology. Increased leaf growth results more often from increased cell expansion rather than increased division. Leaves of crop species exhibit greater increases in leaf thickness than do leaves of wild species. Increased mesophyll and vascular tissue cross-sectional areas, important determinates of photosynthetic rates and assimilate transport capacity, are often reported. Few studies, however, have quantified characteristics more reflective of leaf function such as spatial relationships among chlorenchyma cells (size, orientation, and surface area), intercellular spaces, and conductive tissue. Greater leaf size and/or more leaves per plant are often noted; plants grown in elevated CO<sub>2</sub> exhibited increased leaf area per plant in 66% of studies, compared to 28% of observations reporting no change, and 6% reported a decrease in whole plant leaf area. This resulted in an average net increase in leaf area per plant of 24%. Crop species showed the greatest average increase in whole plant leaf area (+37%) compared to tree species (+14%) and wild, nonwoody species (+15%). Conversely, tree species and wild, nontrees showed the greatest reduction in specific leaf area (–14% and –20%) compared to crop plants (–6%).

Alterations in developmental processes at the shoot apex and within the vascular cambium contributed to increased plant height, altered branching characteristics, and increased stem diameters. The ratio of internode length to node number often increased, but the length and sometimes the number of branches per node was greater, suggesting reduced apical dominance. Data concerning effects of elevated CO<sub>2</sub> on stem/branch anatomy, vital for understanding potential shifts in functional relationships of leaves with stems, roots with stems, and leaves with roots, are too few to

make generalizations. Growth in elevated CO<sub>2</sub> typically leads to increased root length, diameter, and altered branching patterns. Altered branching characteristics in both shoots and roots may impact competitive relationships above and below the ground.

Understanding how increased carbon assimilation affects growth processes (cell division, cell expansion, and cell patterning) will facilitate a better understanding of how plant form will change as atmospheric CO<sub>2</sub> increases. Knowing how basic growth processes respond to increased carbon inputs may also provide a mechanistic basis for the differential phenotypic plasticity exhibited by different plant species/functional types to elevated CO<sub>2</sub>.

*Keywords:* anatomy, development, elevated carbon dioxide, morphogenesis, morphology, ultrastructure

*Received 9 October 1998; resubmitted and accepted 21 December 1998*

## Introduction

Carbon dioxide, released by anthropogenic combustion processes, has increased from 280  $\mu\text{mol mol}^{-1}$  before the onset of the Industrial Revolution to around 365  $\mu\text{mol mol}^{-1}$  today, and continues to rise at about 1.8  $\mu\text{mol mol}^{-1} \text{y}^{-1}$  (Mendelsohn & Rosenberg 1994). Plants exposed to elevated CO<sub>2</sub> often show increased growth and water-use efficiency (Rogers & Dahlman 1993; Allen & Amthor 1995; Wittwer 1995), and increased rates of photosynthesis (Long & Drake 1992; Amthor 1995). Although recent reviews have summarised the effects of elevated CO<sub>2</sub> on ecosystems (Bazzaz 1990; Körner *et al.* 1996), plant herbivore interactions (Bezemer & Jones 1998), crop species (Rogers & Dahlman 1993; Goudriaan & Zadoks 1995), plant biomass accumulation (Poorter 1993), root:shoot ratios (Rogers *et al.* 1997a), and metabolism (Bowes 1991; Farrar & Williams 1991; Long 1991; Stitt 1991; Gunderson & Wulfschleger 1994; Amthor 1995), effects of elevated atmospheric CO<sub>2</sub> on plant structure including ultrastructure, anatomy, morphology, and architecture are far less studied. Changes in plant metabolism resulting from increased C availability will inevitably drive changes in plant structure at multiple hierarchical levels contributing to altered plant, community, and ecosystem level function.

Plant morphogenesis is governed by the effects of environmental conditions superimposed upon genetic constraints. Thus, genetically identical plants can exhibit very different structural features when subjected to different environmental conditions. Although genetically determined design constraints define borders of plasticity, most plant species do mediate the effects of environmental conditions impinging upon them by adjusting developmental processes and resultant structural characteristics to some extent. Ability to adjust both metabolically and structurally to resist stressful environments and to exploit abundant resources will dictate the

fate of individual plant species as the global environment continues to change.

Plants exposed to elevated atmospheric CO<sub>2</sub> are almost always larger than those grown in ambient CO<sub>2</sub>; the magnitude of growth stimulation is typically dependent upon photosynthetic pathway, sink strength, phenotypic plasticity, and plant life history strategies (Hunt *et al.* 1991). Poorter (1993) surveyed the literature (156 plant species) and found the average stimulation of vegetative whole plant growth to be 37%. In addition to increased whole plant biomass, altered root:shoot ratios are often noted, suggesting a shift in the functional relationship between these organs. Rogers *et al.* (1997a) recently reviewed the available literature for crop species and found that root:shoot ratios usually increased (59.5%), sometimes decreased (37.5%), but rarely remained unchanged (3.0%). Although examining plant growth and allocation patterns by assessing total biomass and root:shoot ratios may be a valuable starting point in determining plant response to elevated CO<sub>2</sub>, it may be a rather insensitive indicator of what is actually happening to plant growth in terms of structure and function (Stulen & den Hertog 1993; Taylor *et al.* 1994; Sattler & Rutishauser 1997). For example, differences in root architecture including root depth, branching, and morphology may impact patterns of water and nutrient uptake independent of total biomass (Tremmel & Bazzaz 1993).

The ability of plants to respond to future elevated CO<sub>2</sub> levels will undoubtedly hinge upon physiological characteristics such as sink strength, efficiency of N and water use, and photosynthetic pathway and capacity, but will, with at least equal importance, depend upon plant structural adaptation (Díaz 1995). Körner (1991) has aptly pointed out that the exclusive use of gas exchange data to predict plant success has been over-valued and over-represented in literature addressing plant response to elevated CO<sub>2</sub>. Plant structural responses to elevated CO<sub>2</sub> may prove to be more important than physiological

characteristics in natural environments where plants must compete for scarce resources (Díaz 1995; Teugels *et al.* 1995). For example, Reekie & Bazzaz (1989) reported that although growth in elevated CO<sub>2</sub> did not affect photosynthesis or total biomass accumulation in five trees grown individually, when grown in competition with one another, elevated CO<sub>2</sub> induced changes in species composition resulting from changes in canopy structure. Similarly, Küppers (1985) found that success of species in competition was not related to photosynthetic capacity but instead was related to structural characteristics including branching angles, bud activity, leaf positioning, and internode lengths. Plants must compete for common, usually finite, resources in both natural and managed ecosystems, and these resources are acquired through roots (water and nutrients) and shoots (light, CO<sub>2</sub>). Thus, it follows that in competition, the ability of plants to increase number, size, and efficiency of modules through which resources are acquired, relative to adjacent plants, will define their competitive ability, and resultant success (Teugels *et al.* 1995).

In addition to effects of elevated CO<sub>2</sub> on structurally mediated competitive relationships, alterations in plant form will feed back on physiological processes at the whole plant level which in turn will dictate further growth, development, and survival of the plant. So, as is often expressed by biologists, form and function are inextricably interwoven; function gives form and form results in function. Consideration of physiological alterations, structural modifications, and their interactions resulting from growth in elevated CO<sub>2</sub> will provide a more holistic concept of how plants will change in response to increasing CO<sub>2</sub>, and will help bridge data collected at the physiological level to whole plant and canopy level processes (Murthy & Dougherty 1997).


Plant anatomical, morphological, and architectural adaptations to rising global CO<sub>2</sub> levels may prove to be critical due to the importance of plant form in the acquisition of resources, as a determinate of plant competitive interactions, and as a modifier of metabolic processes. Therefore, the purpose of this manuscript is to review extant data on effects of elevated CO<sub>2</sub> on vegetative plant structure and development. Alterations in plant structural development including ultrastructural, anatomical, morphological, architectural, and canopy levels resulting from exposure to elevated CO<sub>2</sub> will be reviewed and suggestions for further research offered.

### Control of whole plant growth: root shoot signalling

It has long been recognized that plant carbon and N availability are crucial to, and perhaps largely control, whole plant growth patterns. A conceptual understand-

ing of how dynamics of plant C and N pools control growth, via differential allocation of resources to shoots and roots, is central to understanding the significance of rising atmospheric CO<sub>2</sub> for plant structure (Fig. 1). It has long been understood by plant biologists that root-to-shoot ratios of plants as well as whole-plant growth rates, both important determinates of plant success, are determined by the relative availabilities of C and N. In general, plants are able to alleviate disequilibrium between growth processes and resource availability by differentially allocating resources to that plant structure through which the most limiting resource is acquired. For example, plants grown in nutrient-limiting soils allocate resources preferentially below-ground in order to pre-empt critical deficiencies. On the other hand, plants growing in fertile soil are functionally C limited and thus allocate energy into leaf growth to acquire C, the most limiting resource for growth. This concept is reinforced by an extensive literature base (Chapin *et al.* 1987; Aiken & Smucker 1996).

Obtaining a meaningful and satisfying understanding of mechanisms underlying plant structural responses to elevated CO<sub>2</sub> requires a conceptual knowledge of the mechanisms that link recognition (increased photosynthesis → increased carbon → diluted tissue N) with transduction (assimilate partitioning, root-to-shoot signalling, differential gene expression, hormones) with reaction (greater rates of cell division/cell expansion) with adaptation (morphological, anatomical, and ultrastructural). Figure 1 represents a conceptual model illustrating the channels through which increased C input may effect structural change. In some cases, growth responses may be direct (based simply on substrate availability) and significantly more simple than indirect growth responses (based on chemical messengers). Although the complexity of such a holistic view of the interaction of environment with plant growth processes can not be overstated, and perhaps will not be practical for several years, such an approach is vital if form and function are to be linked. In this review, we attempt to understand how plant structure is altered by CO<sub>2</sub>-enriched environments by integrating what is known at scales ranging from molecular to canopy. Closing the gap between individual developmental processes (e.g. cell division, cell expansion, and cell differentiation) and whole plant growth patterns is necessary before we can hope to understand the ways in which plants (both in isolation and within competitive arrays) will respond to future elevated CO<sub>2</sub> levels. As stated recently by Körner *et al.* (1996) concerning the state of knowledge about the implications of rising global CO<sub>2</sub> for plants and ecosystems: '...the field is at the portal of an era where further progress in understanding



**Fig. 1** Conceptual model indicating the channels through which increased atmospheric carbon dioxide availability may effect plant structure. Note the central role of root/shoot meristem function.

CO<sub>2</sub> responses is critically dependent on an effective integration across fields and approaches...'.

## Shoot development

### *Cell division, expansion, and meristem function*

Primary stem growth is ultimately the consequence of apical meristem function. Plant development, initiated at meristems, consists of processes that include cell division, controlled cell expansion, and differentiation (Taylor 1997). These interdependent processes are regulated by specific, genetically programmed, timed ontogenetic events (Körner 1991) integrated with environmental cues to affect the rate, shape and number of organs formed by plants (Kerstetter & Hake 1997). It follows that altered plant structure induced by exposure to elevated CO<sub>2</sub> may be the result of greater rates of cell division, increased cell expansion, altered patterns of primordium initiation, altered morphogenesis and histogenesis, or a combination

of these processes. There has been some disagreement in the literature about which is impacted by elevated levels of CO<sub>2</sub>, thus a brief discussion about factors affecting these processes, in the context of CO<sub>2</sub>-induced changes, is necessary.

First of all, **what stimulates cells to divide?** As recently discussed by Jacobs (1997), there are thought to be two nonmutually exclusive answers to this question. First, cell size homeostasis dictates that as cells expand, a size threshold will be reached beyond which cell volumes and surface areas will exceed the capacity of the nucleus to govern cell function. At this threshold, cells would duplicate their organelles and partition the growing space with new cell walls. Thus, expansion of cells may drive mitosis.

Cells, however, may also divide independently of cell expansion cues. It has long been known that plant hormones including cytokinins, auxins, and gibberellins are involved in controlling developmental events within apical meristems such as cell division, cell

elongation and protein synthesis. For example, both auxins and cytokinins have been shown to increase expression of cyclin genes (Renaudin *et al.* 1994; Kouchi *et al.* 1995; Kende & Zeevaart 1997). It has recently been established in plants that cyclins, a class of regulatory subunits of a family of protein kinases, facilitate the transition of cells from G0 to G1 of the cell cycle, thus stimulating division (Soni *et al.* 1995; Jacobs 1997). It has been reported that sucrose may also be a chemical control point in the cell division cycle (Francis 1992; Ranasinghe & Taylor 1996), perhaps acting by mediating cyclin activity (Kinsman *et al.* 1997). So, cell division may be stimulated by expansion of cells beyond a threshold point, or by increased cyclin activity alone (Jacobs 1997; Kinsman *et al.* 1997). Based on this information, it is possible that growth stimulation of plants grown in elevated CO<sub>2</sub> may be direct (based on substrate supply) or indirect (based on chemical signals) or both. Considering the large impact that growth in elevated CO<sub>2</sub> has on plant root systems (Rogers *et al.* 1997b), it is possible that root cytokinin production, and flux to shoots, may be altered thereby modifying above-ground growth (Fig. 1).

Although it is thought by many researchers that cell production rate largely dictates growth (Körner 1991; Jacobs 1997; Kinsman *et al.* 1997), to a lesser extent increases in **cell expansion** resulting in larger cell size may also contribute to increased plant and organ size without a concomitant increase in cell production (Ranasinghe & Taylor 1996). Cell expansion is controlled by cell wall loosening, wall extensibility, and the rate at which cells can take up water and solutes (Cosgrove 1993, 1997; Ferris & Taylor 1994; Taylor *et al.* 1994). Cell size and rate of cell expansion are influenced by environmental factors including light, water, and nutrient availability and also by endogenous factors such as hormones. It is difficult to disentangle responses of cell production and cell expansion to elevated CO<sub>2</sub> because these processes are so interdependent (Fig. 1). Depending upon context, cell extension may be driven by mitosis, or mitosis may be driven by cell extension (Jacobs 1997) (Fig. 1). This difficulty has been identified and discussed (Ranasinghe & Taylor 1996). Although many studies have reported stimulated growth of stems and branches in plants grown in elevated CO<sub>2</sub> (Downton, Grant & Chacko 1990; Pushnik *et al.* 1995; Slafer & Rawson 1997), few studies have discerned relative contributions of more cell division from larger cell size. St. Omer & Horvath (1984) examined cell size of primary stem tissue for *Layia platyglossa* grown in elevated CO<sub>2</sub>. No increase in diameter of either xylem or sieve elements was evident. Although cell size was unchanged, cortex width and stele diameter of plants grown in 700 µmol mol<sup>-1</sup> CO<sub>2</sub>

were 45 and 41% higher than in plants grown in atmospheres containing 300 µmol mol<sup>-1</sup> CO<sub>2</sub> implying that cell division alone was enhanced.

Kinsman *et al.* (1997) recently provided evidence indicating that exposure to elevated CO<sub>2</sub> stimulates primary growth of shoots by increasing the proportion of rapidly dividing cells and shortening cell cycle durations in shoot apices. Populations of *Dactylus glomerata* from Portugal exhibited a 1.5–3.0 fold increase in the proportion of rapidly cycling cells in the apical dome while a population from Sweden showed only a 1.2 fold increase when exposed to elevated CO<sub>2</sub>. Furthermore, they noted that the cell cycle shortened ≈ 26% in both populations. In the Portuguese population, decreases in the length of the cell cycle resulted almost exclusively from a shortening of the G1 phase, whereas in the Swedish population, both the G1 and G2 phases were shortened. Although the authors did not assay for altered cyclin activity or cytokinin levels or measure tissue carbohydrate levels, they speculated that increased photosynthate availability (i.e. sucrose) in meristems may have increased the proportion of rapidly dividing cells by stimulating cyclin activity. Interestingly, differential increases in total biomass due to exposure to elevated CO<sub>2</sub> paralleled the differential increases in the proportion of dividing cells for the two populations (33 and 21% increases in total biomass for the Portuguese and Swedish populations, respectively). These results suggest that events at the cellular level may provide a key to understanding how growth is controlled at the whole plant level and thereby contribute to a mechanistic understanding of differential phenotypic plasticity exhibited by different species grown at elevated CO<sub>2</sub> concentrations (Körner 1991; Taylor *et al.* 1994). Unfortunately, these results also underscore the problems associated with extending findings observed for one species or population to other related species or populations (Kinsman *et al.* 1996).

### Branching

Although it is clear that exposure of plants to elevated CO<sub>2</sub> stimulates cell division at the shoot apical meristem either directly or indirectly, it is not clear how these cells are partitioned at the shoot axis. Undifferentiated cells produced at the shoot apex must undergo transition to a more specialized state in which they either become components of organ primordia or contribute to internodes between organs (Clark 1997). Examining the effects of CO<sub>2</sub> enrichment on internode length relative to lateral branch or leaf initiation may provide clues for understanding cell partitioning at the shoot apices.

Numerous studies have shown increased **stem or branch elongation** in plants grown in elevated CO<sub>2</sub>

**Table 1** Effects of growth in elevated CO<sub>2</sub> on stem and branch characteristics of woody species

| Species                    | <sup>s</sup> Location / Duration | Ambient [CO <sub>2</sub> ] | Elevated [CO <sub>2</sub> ] | Branch/stem diameter | Branch length | Branch number         | Plant height | Branching patterns   | Anatomical alterations   | Reference                         |
|----------------------------|----------------------------------|----------------------------|-----------------------------|----------------------|---------------|-----------------------|--------------|--|--|-----------------------------------|
| <i>Pinus radiata</i>       | GC (120 d)                       | 320                        | 640                         | -                    | -             | -                     | -            | -  | trach. length NS<br>trach. diameter NS<br>wood density NS        | Donaldson <i>et al.</i> (1987)    |
|                            | GC (22 wks)                      | 330                        | 660                         | -                    | -             | -                     | NS           | -  | wood density +   | Conroy <i>et al.</i> (1986)       |
|                            | GH (2 y)                         | 340                        | 660                         | +                    | -             | +                     | (-)          | whorl # (-)<br>branches/whorl +<br>apical dom. (-)   | trach. length NS<br>trach. wall thickness +44%<br>wood density + | Conroy <i>et al.</i> (1990a)      |
| <i>Pinus taeda</i>         | Phy (1 season)                   | 350                        | 500                         | +                    | -             | NS                    | +            | -  | -  | Sionet <i>et al.</i> (1985)       |
|                            |                                  |                            | 650                         | +                    | -             | NS                    | +            | -  | -  |                                   |
|                            | Phy (113 d)                      | 350                        | 675                         | NS                   | -             | -                     | NS           | -  | -  | Tolley and Strain (1984)          |
|                            |                                  |                            | 1000                        | NS                   | -             | -                     | NS           | -  | -  |                                   |
|                            | <sup>8</sup> BC (21 m)           | 360                        | 535                         | -                    | -16%*         | -                     | -            | flush # NS   | bark density NS  | Murthy and Dougherty (1997)       |
|                            |                                  |                            | 710                         | -                    | +15%          | -                     | -            | flush # NS   | bark density NS  | Larigauderie <i>et al.</i> (1994) |
|                            | Phy (172 d)                      | 375                        | 710                         | LN: +12%             | +47%          | 1°: +45%<br>2°: +66%  | +15%         | plants taller with more<br>and longer<br>1 <sup>st</sup> and 2 <sup>nd</sup> order<br>branches | -  |                                   |
|                            |                                  |                            |                             | HN: +27%             | +53%          | 1°: 1100%<br>2°: +56% | +17%         |  |  |                                   |
| <i>Pinus ponderosa</i>     | GC (6 m)                         | 350                        | 525                         | NS                   | -             | -                     | +20%         | -  | -  | Pushnik <i>et al.</i> (1995)      |
| <i>Picea glauca</i>        | GC (100 d)                       | 350                        | 750                         | -                    | -             | NS                    | NS           | -  | -  | Brown and Higginbotham (1986)     |
| <i>Picea rubens</i>        | GH (5 m)                         | 362                        | 711                         | +33%                 | -             | +33%                  | +23%         | buds yielded<br>greater new<br>fixed growth  | -  | Samuelson and Seiler (1993)       |
| <i>Garcinia mangostana</i> | GC (1 y)                         | 395                        | 800                         | +26%                 | +56%          | +                     | +33%         | 1° node # NA<br>lateral node #<br>+40%   | -  | Downton <i>et al.</i> (1990)      |
| <i>Populus tremuloides</i> | GC (100 d)                       | 350                        | 750                         | -                    | -             | NS                    | NS           | -  | -  | Brown and Higginbotham (1986)     |
| <i>Populus trichocarpa</i> | OTC (92 d)                       | 350                        | 700                         | -                    | +56%          | +                     | +            | -  | -  | Radoglou and Jarvis (1990)        |

| Species                        | <sup>§</sup> Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Branch/stem<br>diameter | Branch length | Branch<br>number | Plant height | Branching patterns   | Anatomical<br>alterations                   | Reference                         |
|--------------------------------|------------------------------------|-------------------------------|--------------------------------|-------------------------|---------------|------------------|--------------|--|---|-----------------------------------|
| <i>Ochroma lagopus</i>         | GC (60 d)                          | 350                           | 675                            | -                       | -             | -                | NS           | -  | -   | Oberbauer <i>et al.</i> (1985)    |
| <i>Pentaclethra macroloba</i>  | GC (123 d)                         | 350                           | 675                            | -                       | -             | -                | NS           | -  | -   |                                   |
| <i>Liquidambar styraciflua</i> | Phy (1 season)                     | 350                           | 500                            | +                       | -             | NS               | +            | NS   | -   | Sionit <i>et al.</i> (1985)       |
|                                | Phy (113 d)                        | 350                           | 650                            | +                       | -             | +122%            | +            | crown shape altered  | -   |                                   |
|                                |                                    |                               | 675                            | +16%                    | -             | -                | +29%         | -  | -   | Tolley and Strain (1984)          |
|                                |                                    |                               | 1000                           | +12%                    | -             | -                | +20%         | -  | -   |                                   |
| <i>Rhizophora mangle</i>       | GH (13 m)                          | 350                           | 700                            | -                       | NS            | +60%             | NS           | branching<br>angles/symmetry NS<br>stem volume +100%<br>URL+ | -   | Farnsworth <i>et al.</i> (1996)   |
| <i>Castanea sativa</i>         | GH (1 season)                      | 350                           | 700                            | LF: NS<br>HF: +18%      | NS            | -                | NS           | NS   | -   | El Kohen <i>et al.</i> (1992)     |
|                                | GC (7 m)                           | 350                           | 700                            | -                       | NS<br>-21%    | -                | NS           | early cessation<br>of stem growth                            | -   | Mousseau and Enoch<br>(1989)      |
| <i>Betula pendula</i>          | GC (79 d)                          | 350                           | 700                            | -                       | -             | NS               | NS           | side shoot #: NS<br>canopy shape NS                          | -   | Pettersson and McDonald<br>(1992) |
| <i>Quercus robur</i>           | GH (19 m)                          | 350                           | 700                            | +175%                   | +137%         | -                | +296%        | -  | increased stem<br>growth rate               | Atkinson <i>et al.</i> (1997)     |
|                                | GH (10 m)                          | 350                           | 700                            | +                       | -             | -                | -            | -  | vessels/stem +<br>vessel lumen              | Atkinson and Taylor<br>(1996)     |
| <i>Prunus avium</i>            | GH (2 m)                           | 350                           | 700                            | -                       | -             | -                | -            | -  | area/vessel +<br>area/stem x-sect.<br>+140% |                                   |
|                                |                                    |                               |                                |                         |               |                  |              |  | vessels/stem NS<br>vessel lumen             |                                   |
|                                |                                    |                               |                                |                         |               |                  |              |  | area/vessel NS                              |                                   |
|                                |                                    |                               |                                |                         |               |                  |              |  | area/stem x-sect. NS                        |                                   |

<sup>§</sup>Plants from all experiments cited in this table were grown in containers except those indicated with a \* (plants grown in the ground). d, days; m, months; y, years; GC, growth chamber; GH, glass house; OTC, open top chamber; phy, phytotron; BC, branch chamber.  
 NS, not significant; LN, low nitrogen; HN, high nitrogen; LF, low fertility; HF, high fertility; \*, mean of two or more fertility or water treatments.  
 Trach, tracheid





without concomitant increases in **node number** (Table 1). For example, Downton *et al.* (1990) reported that *Garcinia mangostana* was 33% taller at elevated CO<sub>2</sub> but that the number of primary nodes was not affected. Rogers *et al.* (1992) reported a 15% increase in plant height for *Glycine max* grown in elevated CO<sub>2</sub> although the number of nodes was unchanged. Similarly, Slafer & Rawson (1997) reported that height of *Triticum aestivum* plants grown in elevated CO<sub>2</sub> increased by 17% resulting from increased internode length, not from greater numbers of nodes. *Lonicera japonica* grown at high CO<sub>2</sub> levels had fewer main stem nodes while plant height was unchanged (Sasek & Strain 1989). Ackerly *et al.* (1992) concluded that increased branch number in *Amaranthus retroflexus* grown in elevated CO<sub>2</sub> was due to effects on overall rate of development, not to changes in pattern of branch initiation. Finally, Sionit *et al.* (1985) observed greater height in *Pinus taeda* grown in elevated CO<sub>2</sub>, but no change in total number of branches. These results suggest that the growth and development of cells and tissues below the site of lateral organ formation are stimulated to a greater extent than is the formation of organ primordia at the shoot tip.


Differential stimulation of cell division in different regions of apical meristems may account for altered patterns of **branch initiation** relative to **internode elongation** (discussed above). Kinsman *et al.* (1996) observed that cell doubling times (cdt) for pith rib meristem cells in *Dactylis glomerata* decreased 4.8 fold at 10 C, 6.1 fold at 20 C, and 2 fold at 30 C in plants grown at 700 µmol mol<sup>-1</sup> CO<sub>2</sub> compared to those grown under ambient CO<sub>2</sub> levels. Cell doubling times in the peripheral meristem zone and central zone (most distal region of apical dome) were not as sensitive to increases in elevated CO<sub>2</sub>. The authors suggested that the pith rib meristem zone would be the first to receive the extra photosynthates produced as a result of the increase in atmospheric CO<sub>2</sub>, but concluded that further investigation is warranted. The central zone is the most distal region of the apical dome and, ultimately, is the source of all cells below it. The peripheral meristem is the location of organ initiation and overall cell patterning, and the pith meristem is the site of cell division and expansion and contributes to axial and radial primary growth of internodes (Esau 1977). Thus, a decrease in the ratio of cdt in the pith rib meristem to cdt in the central zone and the peripheral zone may partly explain why the ratio of internode length to node (organ) number often increases in plants grown in elevated CO<sub>2</sub>. The extent of internodal elongation is perhaps of primary importance in establishing the gross morphology of a species (Esau 1977).

Although node number appears rather insensitive to elevated atmospheric CO<sub>2</sub>, several studies have reported that branch initiation and number have been stimulated,

while plant height or branch length has decreased or remained unchanged (Tables 1 and 2). *Rhizophora mangle* grown in elevated CO<sub>2</sub> had 60% more branches than ambient grown plants, but plant height and branch length were unaffected (Farnsworth *et al.* 1996). They also reported decreased branch plastochron. In *Agrostis capillaris*, growth in elevated CO<sub>2</sub> increased tiller number by 20% while plant height remained unchanged (Newbery & Wolfenden 1996). Similarly, Conroy *et al.* (1990a) reported higher branch numbers, resulting from more branches per whorl, in *Pinus radiata* grown at high CO<sub>2</sub> even though plant height decreased. Several studies have suggested that CO<sub>2</sub> enrichment may increase the number (*Trifolium repens*, Ryle & Powell 1992; *Quercus alba*, Norby *et al.* 1986; *Pinus radiata*, Conroy *et al.* 1990a; *Rhizophora mangle*, Farnsworth *et al.* 1996) or size (*Quercus alba*, Norby *et al.* 1986) of buds/node. Other studies have reported either no effects of elevated CO<sub>2</sub> on branch number and branch/stem length (*Castanea sativa*, Pettersson & McDonald 1992; *Picea glauca*, *Populus tremuloides*, Brown & Higginbotham 1986) or increases in both branch number and branch/stem length (*Populus trichocarpa*, Radoglou & Jarvis 1990a; *Pinus taeda*, Larigauderie *et al.* 1994).

Greater branch elongation and more branches per node resulting from growth in elevated CO<sub>2</sub> may imply **reduced apical dominance** (Tables 1 and 2). Conroy *et al.* (1990a) reported that in *Pinus radiata* grown in elevated CO<sub>2</sub>, the apical portion of the main stem was shorter than the branches at the most terminal whorl which they attributed to reduced apical dominance. Anderson (1976) also observed reduced apical dominance of *Pisum sativum* grown in elevated CO<sub>2</sub>. Reduced apical dominance in plants grown in CO<sub>2</sub> enriched atmospheres may result from altered hormonal production/transport due to effects on apical meristem function, or from alterations in whole plant carbon allocation. Mousseau & Enoch (1989) suggested that reduced branch length (~21%) of *Castanea sativa* induced by CO<sub>2</sub> enrichment was the result of early cessation of stem growth caused by apical bud necrosis. Axillary bud status (domant or active) and lateral branch elongation are not, however, completely controlled or dominated by auxin produced by and transported from apical meristems; overall plant vigor and cues transduced from root stimulation or an improvement in water status, can stimulate lateral growth activity (Stafstrom 1995).

The above discussion shows that exposure of plants to elevated CO<sub>2</sub> may stimulate elongation of branches and stems without accompanying increases in the number of nodes produced. Figure 2 shows architectural diagrams of morphological data obtained during the late pod fill stage of soybean plants grown at [CO<sub>2</sub>] of 349, 645, and 946 ppm under two water regimes, and effectively




**Fig. 2** Diagrams of well-watered (WW) and water-stressed (WS) soybean plants reflecting average morphological data collected at week 14 after planting, later pod fill, from plants grown at 349, 645 and 946 ppm CO<sub>2</sub>. Diagonal lines = the sum of the lateral branch lengths at each node; perpendiculars = leaves; dots = 2 pods; scale = 10 cm; *N* = 6 (Prior 1986).

illustrates how branch length is stimulated relative to branch number (Prior 1986). It is important to note however, that although the ratio of height/length to node number appears to increase in plants grown in elevated CO<sub>2</sub>, branch number often increases, resulting mainly from greater absolute numbers of nodes per plant, or less frequently, from more buds per node. These alterations, indicating shifts in shoot allometry, have been reported to change canopy shape (*Lonicera japonica*, Sasek & Strain 1991) perhaps contributing to altered competitive relationships between different plant species (*Liquidambar styraciflua* vs. *Pinus taeda*, Sionit *et al.* 1985; Reekie & Bazzaz 1989). Reekie & Bazzaz (1989) grew tropical tree species including *Cecropia obtusifolia*, *Myriocarpa longipes*, *Piper auritum*, *Senna multijuga*, and *Trichospermum mexicanum* in competition at 350, 525, or 700 ppm CO<sub>2</sub>. They found that *Senna* decreased in importance resulting from a reduction in the height at which leaves were displayed (Fig. 3). In contrast, *Trichospermum*, *Piper* and *Cecropia* all increased in importance since their leaves were displayed higher (Fig. 3). The authors concluded that shifts in community composition were brought about by alterations in canopy structure due to changes in height growth and branching patterns (Reekie 1996), caused ultimately by alterations in developmental processes within meristematic tissue. Changes in primary stem growth need to be examined and compared for multiple species if we hope to be able to make useful generalizations (based on growth form or functional type) about how CO<sub>2</sub> enrichment affects branching, canopy characteristics, and competitive ability.

#### Secondary tissues and anatomy

Below the shoot apex, at the shoot or branch site where elongation (primary growth) has ceased, secondary growth, resulting in the production of secondary xylem (wood) and phloem (bark), occurs by activation of a vascular cambium (Esau 1977). Although most studies concerned with plant structural responses to their environment have focused on leaves, the importance of secondary stem tissue in mediating the flux of resources acquired above and below the ground can not be overstated. In the words of Kozlowski & Pallardy (1997), 'Transport of adequate supplies of metabolites to meristematic tissues where they are used as respiratory subunits and building materials is as important for plant growth as the physiological processes that produce them'. Stem characteristics including anatomical and ultrastructural characteristics of the individual cells which comprise functional domains, in addition to spatial arrangements of the cells and domains that together comprise functional phloem and xylem networks, are crucial. Stems are perhaps the least able to



**Fig. 3** Effects of atmospheric CO<sub>2</sub> concentration on leaf area profiles of tropical trees. C = *Cecropia*, M = *myriocarpa*, P = *Piper*, S = *Senna*, T = *Trichospermum*. Each tick on the horizontal axis = 0.01 m<sup>2</sup>; leaf area was summed over 10 cm intervals. Numbers above each species represent mean canopy height (m). From Reekie & Bazzaz (1989), with permission from Springer-Verlag.

respond to differential atmospheric and edaphic conditions because they grow and are maintained using resources which are all acquired elsewhere; stems must extract required substances through the conduits which pass through them (Cheeseman *et al.* 1996). As such, understanding how stem, leaf and root growth are coordinated represents a complicated problem.

Increases in **diameter of stems and branches** have been reported for many species growing under conditions of elevated CO<sub>2</sub> (Tables 1 and 2) (*Garcinia mangostana*, Downton *et al.* 1990; *Liquidambar styraciflua*, *Pinus taeda*, Sionit *et al.* 1985; *Castanea sativa*, Samuelson & Seiler 1993; *Quercus robur*, Atkinson *et al.* 1997; *Pinus radiata*,

**Table 3** Effects of growth in elevated CO<sub>2</sub> on leaf characteristics of tree species

| Species                                    | <sup>s</sup> Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf | Leaf<br>area/plant   | Leaf<br>#/plant | SLA   | Other leaf<br>changes         | Reference                         |
|--|------------------------------------|-------------------------------|--------------------------------|-----------|----------------------|-----------------|-------|-------------------------------|-----------------------------------|
| <i>Pinus ponderosa</i>                     | GC (6 m)                           | 350                           | 525                            | -         | -33%                 | -               | -14%  | mes. area -8%<br>vt. area +4% | Pushnik <i>et al.</i> (1995)      |
| <i>Pinus taeda</i>                         | GC (6 m)                           | "                             | 700                            | -         | -47%                 | -               | -14%  | leaf thickness +              | Rogers <i>et al.</i> (1983)       |
|  | OTC (3 m)                          | 340                           | 520                            | -         | -                    | -               | -     | " ++                          |                                   |
|  |                                    |                               | 718                            | -         | -                    | -               | -     | " +110%                       | Tolley & Strain (1984)            |
|  | Phy (113 d)                        | 350                           | 675                            | -         | NS                   | -               | -9%   | Leaf area duration NS         |                                   |
|  | Phy (113 d)                        | "                             | 1000                           | -         | NS                   | -               | -2%   | needle dry wt: +58%           | Larigauderie <i>et al.</i> (1994) |
|  | Phy (172 d)                        | 375                           | 710                            | -         | LN: +47%<br>HN: +66% | -               | -8%   | " +71%                        |                                   |
| <i>Pinus radiata</i>                       | gBC (21 m)                         | 360                           | 535                            | -         | NS                   | NS              | NS    | leaf thickness NS             | Murthy & Dougherty (1997)         |
|  |                                    |                               | 710                            | -         | +16%                 | +12%            | NS    | leaf length NS                |                                   |
|  | OTC (45 d)                         | 340                           | 520                            | -         | -                    | -               | -     | leaf thickness NS             | Thomas & Harvey (1983)            |
|  |                                    |                               |                                |           |                      |                 |       | mes. area NS                  |                                   |
|  |                                    |                               |                                |           |                      |                 |       | e-ft-vt area NS               |                                   |
|  |                                    |                               | 718                            | -         | -                    | -               | -     | pid. area (x-sect) NS         |                                   |
|  |                                    |                               |                                |           |                      |                 |       | leaf thickness NS             |                                   |
|  |                                    |                               |                                |           |                      |                 |       | mes. area NS                  |                                   |
|  |                                    |                               |                                |           |                      |                 |       | e-ft-vt area +8%              |                                   |
|  |                                    |                               | 910                            | -         | -                    | -               | -     | epid. area (x-sect) NS        | Conroy <i>et al.</i> (1986)       |
| <i>Pinus palustris</i>                     | GC (22 weeks)                      | 330                           | 660                            | +15%*     | +50%*                | +31%*           | -16%* | mes. area +10%                |                                   |
|  |                                    |                               |                                |           |                      |                 |       | leaf thickness +10%           |                                   |
|  |                                    |                               |                                |           |                      |                 |       | e-ft-vt area +11%             |                                   |
|  |                                    |                               |                                |           |                      |                 |       | epid. area (x-sect) NS        |                                   |
|  |                                    |                               |                                |           |                      |                 |       | low P: t-tissue -14%          | Conroy <i>et al.</i> (1990a)      |
|  |                                    |                               |                                |           |                      |                 |       | high P: " +23%                |                                   |
|  | GH (2 years)<br>OTC (20 m)         | 340<br>360                    | 660<br>720                     | -         | NS                   | NS              | -     | low P: mes. area NS           | Pritchard <i>et al.</i> (1997)    |
|  |                                    |                               |                                |           |                      |                 |       | high P: ' +38%                |                                   |
|  |                                    |                               |                                |           |                      |                 |       | low P: vt. area NS            |                                   |
|  |                                    |                               |                                |           |                      |                 |       | high P: ' NS                  |                                   |
| <i>Picea glauca</i><br><i>Picea rubens</i> | OTC (12 m)                         | 360                           | 720                            | -         | -                    | -               | -     | Phloem area -25%              | Pritchard <i>et al.</i> (1998)    |
|  |                                    |                               |                                |           |                      |                 |       | # sieve cells -26%            |                                   |
|  |                                    |                               |                                |           |                      |                 |       | sieve cell size NS            |                                   |
|  |                                    |                               |                                |           |                      |                 |       | leaf x-sect. area +15%        |                                   |
|  |                                    |                               |                                |           |                      |                 |       | fascicle volume +8%           |                                   |
|  | " (20 m)                           |                               |                                | -         | -                    | -               | -     | t-tis area +3%                |                                   |
|  |                                    |                               |                                |           |                      |                 |       | mes. area +17%¶               |                                   |
|  |                                    |                               |                                |           |                      |                 |       | vt. area +7%¶                 |                                   |
|  |                                    |                               |                                |           |                      |                 |       | needle length NS              |                                   |
|  |                                    |                               |                                |           |                      |                 |       | leaf x-sect. area NS          |                                   |
|  |                                    |                               |                                |           |                      |                 |       | fascicle volume -8%           |                                   |
|  |                                    |                               |                                |           |                      |                 |       | t-tis area NS                 |                                   |
|  |                                    |                               |                                |           |                      |                 |       | mes. area -17%¶               |                                   |
|  |                                    |                               |                                |           |                      |                 |       | vt. area -10%¶                |                                   |
|  |                                    |                               |                                |           |                      |                 |       | needle length -7%¶            |                                   |
| <i>Picea glauca</i>                        | GC (100 d)                         | 350                           | 750                            | -         | NA                   | -               | -15%* | leaf wt. ratio NS             | Brown and Higginbotham (1986)     |
| <i>Picea rubens</i>                        | GH (5 m)                           | 362                           | 711                            | -         | -                    | -               | -22%* | -                             | Samuelson & Seiler (1993)         |

| Species                        | <sup>§</sup> Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf       | Leaf<br>area/plant | Leaf<br>#/plant | SLA          | Other leaf<br>changes                            | Reference                       |
|--------------------------------|------------------------------------|-------------------------------|--------------------------------|-----------------|--------------------|-----------------|--------------|--|---------------------------------|
| <i>Garcinia mangostana</i>     | GC (1 years)                       | 395                           | 800                            | +10%            | +28%               | +18%            | -19%         | -  | Downton <i>et al.</i> (1990)    |
| <i>Liquidambar styraciflua</i> | OTC (3 m)                          | 340                           | 520                            | -               | -                  | -               | -            | leaf thickness +<br>"++"                         | Rogers <i>et al.</i> (1983)     |
|                                |                                    |                               | 718                            | -               | -                  | -               | -            | "++"   |                                 |
|                                | Phy (113 d)                        | 350                           | 675                            | NS              | NS                 | +               | -16%         | leaf area duration +56%                          | Tolley & Strain (1984)          |
|                                |                                    |                               | 1000                           | NS              | NS                 | +               | -28%         | leaf area duration +22%                          |                                 |
|                                | OTC (45 d)                         | 340                           | 520                            | -               | -                  | -               | -            | leaf thickness +17%                              | Thomas & Harvey (1983)          |
|                                |                                    |                               |                                |                 |                    |                 |              | spongy mes. NS                                   |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | palisade mes. +24%                               |                                 |
|                                |                                    |                               | 718                            | -               | -                  | -               | -            | epid. area (x-sect) NS                           |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | leaf thickness +25%                              |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | spongy mes. +21%                                 |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | palisade mes. +36%                               |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | epid. area (x-sect) NS                           |                                 |
|                                |                                    |                               | 910                            | -               | -                  | -               | -            | leaf thickness +21%                              |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | spongy mes. +14%                                 |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | palisade mes. +33%                               |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | epid. area (x-sect) NS                           |                                 |
| <i>Quercus alba</i>            | GC (24 weeks)                      | 389                           | 496                            | -               | NS                 | -               | -            | -  | Norby & O'Neill (1989)          |
|                                |                                    |                               | 793                            | -               | NS                 | -               | -            | -  |                                 |
| <i>Populus</i> spp.            | GC (100 d)                         | 350                           | 750                            | -               | NS                 | -               | -15%*        | leaf wt. ratio NS                                | Brown and Higginbotham (1986)   |
|                                | gOTC (158 d)                       | 345                           | 693                            | LN: NS<br>HN: + | NS<br>+35%         | NS<br>++        | -12%<br>-16% | leaf area duration NS<br>leaf area duration +38% | Curtis <i>et al.</i> (1995)     |
|                                | OTC (92 d)                         | 350                           | 700                            | NS              | +39%               | +11%            | (-)          | leaf thickness +13%                              | Radoglou & Jarvis (1990a)       |
|                                |                                    |                               |                                |                 |                    |                 |              | spongy mes. (+)                                  |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | palisade mes. NS                                 |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | epid. area (x-sect) +<br>cell size NS            |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | rate of growth +                                 |                                 |
| <i>Castanea sativa</i>         | GH (1 season)                      | 350                           | 700                            | -               | LF: NS<br>HF: +24% | -               | NS           | -  | El Kohen <i>et al.</i> (1992)   |
|                                |                                    |                               |                                |                 |                    |                 | NS           | -  |                                 |
|                                | GC (1 season)                      | 360                           | 700                            | LF: NS          | NS                 | NS              | -            | -  | El Kohen & Mousseau (1994)      |
|                                |                                    |                               |                                |                 | HF: +24%           | +               | -            | -  |                                 |
| <i>Betula pendula</i>          | GC (70 d)                          | 350                           | 700                            | -               | -                  | NS              | -12%         | -  | Pettersson & McDonald (1992)    |
| <i>Rhizophora mangle</i>       | GH (13 m)                          | 350                           | 700                            | NS              | +30%               | +15%            | NS           | mes. area NS                                     | Farnsworth <i>et al.</i> (1996) |
|                                |                                    |                               |                                |                 |                    |                 |              | epid. area (x-sect) +<br>epid. cell size +       |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | #leaves/branch NS                                |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | leaf area duration NS                            |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | rate of growth +                                 |                                 |
|                                |                                    |                               |                                |                 |                    |                 |              | leaf thickness NS                                |                                 |
| <i>Ochroma lagopus</i>         | GC (60 d)                          | 350                           | 675                            | -               | +39%               | -               | -38%         | leaf area ratio -22%                             | Oberbauer <i>et al.</i> (1985)  |
| <i>Pentaclethra macroloba</i>  | GC (123 d)                         | 350                           | 675                            | -               | NS                 | -               | -20%         | leaf thickness NS<br>leaf area ratio -18%        |                                 |

<sup>§</sup>Plants from all experiments cited in this table were grown in containers unless indicated with a <sup>§</sup> (plants grown in the ground). d, days; m, months; y, years.

GC, growth chamber; GH, glass house; OTC, open top chamber; phy, phytotron; BC, branch chamber; NS, not significant; LN, low nitrogen; HN, high nitrogen; LF, low fertility; HF, high fertility; <sup>¶</sup>, not statistically significant; \*, mean of fertility or water treatments; mes., mesophyll tissue; vt., vascular tissue; e-tt-vt, endodermis-transfusion tissue-vascular tissue; epid, epidermis; t-tis., transfusion tissue; low P, low phosphorus availability; high P, high phosphorus availability. SLA, specific leaf area (leaf area/total leaf dry weight)

Conroy *et al.* 1990a). A few studies have reported that exposure to elevated CO<sub>2</sub> resulted in no effect on secondary growth (i.e. diameter) of stems (*Pinus ponderosa*, Pushnik *et al.* 1995; *Pinus taeda*, Tolley & Strain 1984); and, to our knowledge, no species has exhibited decreased secondary growth in elevated CO<sub>2</sub>. Further evidence suggesting an effect of elevated CO<sub>2</sub> on secondary growth of woody stems is provided from tree ring data (reviewed in Weber & Grulke 1995).

There is a paucity of studies on **stem anatomical characteristics** (Tables 1 and 2). Obtaining an understanding of the functional relationship of leaves and stems, and roots and stems, will be impossible until more information on anatomical alterations in stems is available. For example, stem cross-sectional area is thought to be related to whole plant reproductive capacity (Atkinson & Taylor 1996) and is directly proportional to the amount of leaf area which must be supplied with water and solutes (Zimmermann 1983; Atkinson & Taylor 1996). Anatomical characteristics of individual cells within stems are also very important. The ability of stems to transport water to maintain favourable whole plant water relations is governed not only by quantity, but also by the size of xylem conduits (Tyree & Alexander 1993; Atkinson & Taylor 1996).

Tyree & Alexander (1993) hypothesized that increased photosynthesis, increased carbohydrate availability, and resultant increases in the **numbers and sizes of xylem elements** (tracheids and vessel members) within plants grown in elevated CO<sub>2</sub> may enhance the risk of xylem cavitation. Atkinson & Taylor (1996) examined stem characteristics of *Quercus robur* and *Prunus arium* grown under CO<sub>2</sub> enrichment. For *Quercus*, there were greater numbers of vessels in the mid-stem area contributing to a 140% increase in total stem vessel lumen area. Additionally, they found that xylem elements from *Quercus* plants grown in elevated CO<sub>2</sub> had significantly greater mean vessel lumen area than vessels from plants grown with ambient air. These changes contributed to increased hydraulic conductance in plants grown in elevated compared to ambient CO<sub>2</sub>. It may also be important to note that increases in total stem vessel area were not related to increases in leaf area. This suggests an uncoupling of the functional relationship of leaves with stems which may imply altered predisposition to xylem cavitation due to drought or freezing stress (Atkinson & Taylor 1996). In contrast, *Prunus* stem total vessel lumen area, vessel number, and lumen area per vessel were not affected by elevated CO<sub>2</sub>. Similarly, Donaldson *et al.* (1987) found that CO<sub>2</sub> enrichment caused no differences in tracheid length, tracheid lumen diameter, or tracheid wall thickness in *Pinus radiata*. Conversely, Conroy *et al.* (1990a) reported tracheid wall thickness to increase by 44% in *Pinus radiata* grown under CO<sub>2</sub> enrichment. The

authors suggested these inconsistent results were brought about by dissimilar source-sink relationships due to differences in plant age between the two studies. This type of discrepancy is not unusual in the literature on effects of elevated CO<sub>2</sub> on plant processes, both metabolic and structural. Often effects of elevated CO<sub>2</sub> show subtle differences if growth conditions change: pot size (Arp 1991), nutrient and water availability (Prior *et al.* 1997), and different CO<sub>2</sub> exposure systems and concentrations (Ceulemans & Mousseau 1994). Furthermore, whole plant responses to elevated CO<sub>2</sub> may decrease over time due to biochemical (e.g. decreased rubisco activity, Gunderson & Wullschlegler 1994), ultrastructural (e.g. chloroplast disruption, Pritchard *et al.* 1997), or canopy level (e.g. self-shading, Newbery & Wolfenden 1996) limitations. There are too few data concerning the effects of CO<sub>2</sub> enrichment on secondary stem anatomy to make generalizations.

In addition to increased stem diameter and altered stem cell sizes, **stem density** may be altered by CO<sub>2</sub> enrichment (Tables 1 and 2). Wood density has been reported to increase (*Liquidambar styraciflua*, Rogers *et al.* 1983; *Pinus radiata*, Conroy *et al.* 1986) or remain unchanged by growth in elevated CO<sub>2</sub> (*Pinus taeda*, Murthy & Dougherty 1997). In the only known study to examine density of bark, Murthy & Dougherty (1997) reported no change in bark density of *Pinus taeda* grown in elevated CO<sub>2</sub>. Density of wood may be an important component of wood quality for both timber and paper production (Conroy *et al.* 1990a,b).

**In conclusion**, existing data suggest that elevated CO<sub>2</sub> drives increased stem growth primarily by stimulating cell division within shoot apices. However, cell proliferation may be stimulated to different extents throughout different meristematic regions which perhaps accounts for observed increases in nodal elongation relative to branch initiation, and other shifts in whole plant architecture. Furthermore, in some cases, growth in elevated CO<sub>2</sub> appears to stimulate lateral growth suggesting reduced apical dominance. Although it can be inferred that vascular cambium activity is stimulated (increased secondary stem diameters are often observed), there is a profound lack of data concerning the impact of elevated CO<sub>2</sub> on processes driving secondary growth and also on anatomical shifts which may result from altered cambium function.

## Leaf development

### *Why study effects of elevated CO<sub>2</sub> on leaf structure?*

Of all plant organs, leaves are the most morphologically diverse (Poethig 1997) and exhibit the greatest structural plasticity in response to disparate environmental condi-

tions (Esau 1977). Leaf development is crucial to plant function since leaves are vital to light interception, photosynthesis, water use, and therefore, total plant productivity (Teskey *et al.* 1987; Murthy & Dougherty 1997). Leaf structural adaptations clearly play a central role in adaptation by plants to changing environments (Lewis 1972; Ticha 1982; Ashton & Berlyn 1994). Functionally, developing leaves are thought to also produce a hormonal signal (i.e. auxin) which stimulates differentiation of xylem. Thus, amount of leaf area dictates stem area produced (Taylor *et al.* 1994; Atkinson & Taylor 1996), assuring functional equilibrium between leaves and stems. In a broader context, rates of leaf development, leaf area duration, and leaf efficiency (a function of anatomy, ultrastructure, and biochemistry) throughout the growing season dictate the rate of canopy closure and the yearly canopy productivity index (CPI, annual production of wood per unit of leaf area). A thorough understanding of how increasing CO<sub>2</sub> will impact the dynamics of leaf initiation, morphogenesis (development of shape), histogenesis (development of internal organization) and phenology will be necessary to: (i) determine the impact of elevated CO<sub>2</sub> on leaf and whole plant function (ii) determine the effects of these functional shifts on ecosystem processes and physiognomy, and (ii) more accurately link vegetation processes with global carbon models and budgets.

#### *Leaf initiation*

Leaves usually arise from the surface of meristems as primordial ridges or flattened bumps at predictable intervals (plastochrons) and locations (phyllotaxy) around the axis of the plant (Poethig 1997). Though generally associated with the shoot apical meristem, leaves may arise elsewhere (Clark 1997; Poethig 1997). As recently discussed by Poethig (1997), the relationship between the shoot apical meristem and initiation of leaves has never been established; 'the meristem may represent a region (or type of tissue) in which leaves can spontaneously self-organize rather than a structural entity that makes leaf primordia.' Thus, leaf morphogenesis is discussed separately from shoot morphogenesis in order to reflect this functional discontinuity.

Two models, discussed by Poethig (1997), explain how **leaf primordia** may arise. In the first, it is hypothesized that a field of cells emerges in the meristem having circularly arranged cellulose microfibrils. Such a microfibril arrangement would cause these cells to expand out of the plane of the shoot apical meristem forming the primordial leaf. The second model suggests that leaf primordia may arise as a result of buckling in outer layers of the meristem in response to mechanical stress in

the shoot apex caused by an excess of tissue unable to expand laterally. Such biophysical models, in which physical tension across the meristem triggers changes in cell division patterns leading to the initiation of organ primordia, have recently received molecular support (Taylor 1997).

Plants grown in elevated CO<sub>2</sub> typically have increased rates of photosynthesis leading to greater carbohydrate availability (Cave *et al.* 1981). Increases in assimilate transport and carbohydrate availability in shoot apical meristems may increase rates of cell division as discussed above (Kinsman *et al.* 1997). Furthermore, increased tissue water availability resulting from enhanced water-use efficiency (WUE) and increased root proliferation could contribute to greater rates of cell expansion due to increased cell turgor pressure (Sasek & Strain 1989). These factors combined intuitively suggest increased growth, excess tissue, mechanical stress, increased buckling within apical meristems, and therefore increased leaf initiation in plants exposed to elevated CO<sub>2</sub> concentrations. However, as discussed earlier, cell division may be stimulated to a greater extent in pith rib meristems than in the peripheral meristem region (the site of lateral organ initiation).

Although it is difficult to determine from the literature what effect elevated CO<sub>2</sub> has on leaf initiation at apical meristems, it is generally thought that exposure of plants to elevated CO<sub>2</sub> has only small, if any, effects on **rates of leaf initiation per se** (Ackerly *et al.* 1992). Studies which provide clues about initiation of leaf primordia relative to stem or branch growth are rare, so it is difficult to determine whether increases in whole plant leaf area result from altered allometric relationships between total stem/branch length and leaf number, or if leaf area simply increases in proportion to greater total branch/stem length. Of 10 reports on leaf area ratio (LAR=total projected leaf area/total plant dry weight), 50% reported decreases, 30% increases, and 20% no effect, resulting in an average net reduction on LAR of 16% (Tables 3, 4, and 5). Summarizing the findings for 20 observations of leaf weight ratio (LWR=leaf weight/total plant weight), decreases were observed for 45% of species, no change for 55% of species, and LWR never increased. Average net decrease in LWR was 10%. Decreases in LAR and LWR suggest that plants allocate less carbon to production of new leaf area at elevated levels of CO<sub>2</sub>. So, although faster rates of leaf initiation are sometimes reported, these increases are probably not of the same magnitude as are increases in stem and root growth. Norby (1996) reported the average increase in the canopy productivity index to be 29% in seven tree species grown in elevated CO<sub>2</sub> (650–700 ppm) further suggesting that less leaf area is

**Table 4** Effects of growth in elevated CO<sub>2</sub> on leaf characteristics of crop species

| Species                     | <sup>s</sup> Location/<br>Duration                             | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf                          | Leaf<br>area/plant | Leaf<br>#/plant | SLA  | Other leaf<br>changes  | Reference                     |
|-----------------------------|--|-------------------------------|--------------------------------|------------------------------------|--------------------|-----------------|------|--|-------------------------------|
| <i>Phaseolus vulgaris</i>   | GC (25 d)  | 340                           | 640                            | +                                  | -                  | -               | -    | rate of growth +<br>epid. cell size +<br>XET activity +  | Ranisinghe and Taylor (1996)  |
|                             | GC   | 400                           | 1200                           | -                                  | +26%               | -               | -    | -  | O'Leary and Knecht (1981)     |
|                             | OTC (30 d)   | 350                           | 700                            | -                                  | LF: +20%           | -               | -    | leaf thickness +<br>spongy mes. +<br>palisade mes. +<br>cell # NS<br>cell size +<br>mes. air space (-) | Radoglou and Jarvis (1992)    |
|                             |  |                               |                                |                                    | HF: +19%           | -               | -    | leaf thickness +<br>spongy mes. +<br>palisade mes +<br>cell # NS<br>cell size +<br>mes. air space (-)  |                               |
| <i>Lolium perenne</i>       | GH (1 season)  | 360                           | 710                            | -                                  | +                  | -               | -    | leaf inclination NS<br>leaf distribution NS  | Teugels <i>et al.</i> (1995)  |
|                             | GC (5 wks)   | 340                           | 680                            | NS                                 | -                  | -               | -    | leaf length NS<br>leaf width NS<br>epid. cell length NS  | Ryle and Stanley (1992)       |
|                             | OTC (35 d)   | 371                           | 700                            | aT: +80%<br>a+4°C: +14%<br>aT -15% | -                  | -               | -    | epid. cell density NS<br>mes. area +46%*   | Ferris <i>et al.</i> (1996)   |
|                             | spring leaves<br>OTC (35 d)<br>summer leaves<br>Phy (1 season) | 371                           | 700                            | a+4°C: -13%                        | -                  | -               | -    | epid. area +39%*<br>mes. area -29%*<br>epid. area -14%*  |                               |
| <i>Festuca arundinaceae</i> | GH (1 season)  | 340                           | 680                            | -                                  | -                  | -               | -27% | leaf weight ratio NS   | Morison and Gifford (1984a,b) |
|                             |  | 360                           | 700                            | -                                  | +                  | -               | -    | leaf inclination NS<br>leaf distribution NS  | Teugels <i>et al.</i> (1995)  |
| <i>Triticum aestivum</i>    | GH (1 season)  | 360                           | 720                            | -                                  | -                  | NS              | -    | leaf development NS<br>spikelets/spike NS<br>leaf weight ratio NS                                      | Slafer and Rawson (1997)      |
| <i>Trifolium repens</i>     | Phy (1 season)   | 340                           | 680                            | -                                  | +68%               | -               | -18% | -  | Morison and Gifford (1984a,b) |
|                             | GH (2 m)   | 350                           | 700                            | +                                  | +                  | +               | +    | -  | Jongen <i>et al.</i> (1996)   |
|                             | Phy (1 season)   | 340                           | 680                            | -                                  | +39%               | -               | -14% | leaf weight ratio -14%   | Morison and Gifford (1984a,b) |
|                             | CC (90 d)  | 340                           | 680                            | +5%                                | +                  | +30%            | -10% | leaves/stolon NS<br>stolons/plant +44%   | Ryle and Powell (1992)        |



| Species                        | §Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf | Leaf<br>area/plant | Leaf<br>#/plant | SLA  | Other leaf<br>changes   | Reference                     |
|--------------------------------|------------------------|-------------------------------|--------------------------------|-----------|--------------------|-----------------|------|---|-------------------------------|
| <i>Lycopersicon esculentum</i> | GH (29 d)              | 400                           | 1000                           | -         | -                  | -               | -    | leaf thickness +<br>cell size +<br>cell # NS                          | Madsen (1968)                 |
| <i>Gossypium hirsutum</i>      | GH (40 d)              | 330                           | 640                            | -         | +60%               | -               | -    | -   | Wong (1979)                   |
|                                | Phy (1 season)         | 340                           | 680                            | -         | +14%               | -               | -18% | leaf weight ratio NS  | Morison and Gifford (1984a,b) |
| <i>Zea mays</i>                | GH (40 d)              | 330                           | 640                            | -         | +10%               | -               | -    | -   | Wong (1979)                   |
|                                | OTC (3 m)              | 340                           | 520                            | -         | +                  | -               | -    | leaf thickness NS   | Rogers <i>et al.</i> (1983)   |
|                                |                        |                               | 718                            | -         | ++                 | -               | -    | -   |                               |
|                                |                        |                               | 910                            | -         | +++                | -               | -    | -   |                               |
|                                | OTC (45 d)             | 340                           | 520                            | -         | -                  | -               | -    | leaf thickness NS<br>mes. area -11%<br>epid. area (x-sect) NS         | Thomas and Harvey (1983)      |
|                                |                        |                               | 718                            | -         | -                  | -               | -    | leaf thickness -9%<br>mes. area -11%<br>epid. area (x-sect) NS        |                               |
|                                |                        |                               | 910                            | -         | -                  | -               | -    | leaf thickness NS<br>mes. area NS                                     |                               |
| <i>Helianthus annuus</i>       | Phy (1 season)         | 340                           | 680                            | -         | +40%               | -               | NS   | epid. area (x-sect) NA<br>leaf weight ratio -13%                      | Morison and Gifford (1984a,b) |
| <i>Vigna unguiculata</i>       | Phy                    | 340                           | 680                            | -         | +14%               | -               | -18% | leaf weight ratio -17%  | Morison and Gifford (1984a,b) |
| <i>Macroptilium purpureum</i>  | Phy                    |                               |                                | -         | +56%               | -               | -22% | leaf weight ratio -14%  |                               |
|                                | Phy                    |                               |                                | -         | +39%               | -               | NS   | leaf weight ratio NS  |                               |
| <i>Oryza sativa</i>            | Phy                    |                               |                                | -         | NS                 | -               | -14% | leaf weight ratio NS  |                               |
| <i>Phalaris aquatica</i>       | Phy                    |                               |                                | -         | +31%               | -               | NS   | leaf weight ratio NS  |                               |
| <i>Amaranthus</i> spp.         | Phy                    | 340                           | 680                            | -         | +15%               | -               | -11% | leaf weight ratio -15%  | Ackerly <i>et al.</i> (1992)  |
|                                | GH (31 d)              | 400                           | 700                            | -         | @28°C: +93%        | -               | -    | plastchr. index +19%<br>stem leaf area +43%<br>branch leaf area +180% |                               |
|                                |                        |                               |                                | -         | @38°C: -75%        | -               | -    | plastchr. index NS<br>stem leaf area NS                               |                               |
| <i>Sorghum bicolor</i>         | GH (1 season)          | 350                           | 700                            | -         | (-)                | -               | (-)  | branch leaf area -68%<br>leaf area ratio NS                           | Carbutt <i>et al.</i> (1990)  |
| <i>Brassica napus</i>          | Phy (1 season)         | 340                           | 680                            | -         | +29%               | -               | NS   | leaf weight ratio -15%  | Morison and Gifford (1984a,b) |
| <i>Hordium vulgare</i>         | Phy "                  | "                             | "                              | -         | +40%               | -               | -30% | leaf weight ratio NS  |                               |
| <i>Vicia faba</i>              | Phy "                  | "                             | "                              | -         | +57%               | -               | -16% | leaf weight ratio NS  |                               |
|                                | Phy "                  | "                             | "                              | -         | +42%               | -               | -11% | leaf weight ratio NS  |                               |

Table 4. Continued

| Species                | <sup>s</sup> Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf | Leaf<br>area/plant | Leaf<br>#/plant | SLA  | Other leaf<br>changes   | Reference   |
|------------------------|------------------------------------|-------------------------------|--------------------------------|-----------|--------------------|-----------------|------|---|---|
| <i>Pisum sativum</i>   | Phy "                              | "                             | "                              | -         | +53%               | -               | -10% | leaf weight ratio NS  | Leadley <i>et al.</i> (1987)                            |
| <i>Medicago sativa</i> | Phy "                              | "                             | "                              | -         | +75%               | -               | -44% | leaf weight ratio NS  |   |
| <i>Raphanus sativa</i> | Phy "                              | "                             | "                              | -         | -                  | -               | -14% | leaf weight ratio -41%  |   |
| <i>Glycine max</i>     | OTC (45 d)                         | 348                           | 645                            | -         | -                  | -               | -    | leaf thickness +<br>vt area:total leaf area +<br>mes. cell surface area (-) |   |
|                        | OTC (3 m)                          | 340                           | 520                            | -         | -                  | -               | -    | leaf thickness +<br>palisade mes. area +                                    | Rogers <i>et al.</i> (1983)                             |
|                        |                                    |                               | 718                            | -         | -                  | -               | -    | leaf thickness ++   |   |
|                        |                                    |                               | 910                            | -         | -                  | -               | -    | palisade mes. area +  |   |
|                        |                                    |                               | 700                            | +56%      | -                  | -               | -    | leaf thickness +131%  |   |
|                        | Phy (18 d)                         | 350                           |                                |           |                    |                 |      | leaf x-sect. area +12%  | Rogers <i>et al.</i> (1992)<br>Thomas and Harvey (1983) |
|                        | OTC (45 d)                         | 340                           | 520                            | -         | -                  | -               | -    | leaf thickness +35%   |   |
|                        |                                    |                               |                                |           |                    |                 |      | palisade mes. area +32%<br>spongy mes. area +45% <sup>†</sup>               |   |
|                        |                                    |                               |                                |           |                    |                 |      | epid. area (x-sect) NS  |   |
|                        |                                    |                               | 718                            | -         | -                  | -               | -    | leaf thickness +33%   | Vu <i>et al.</i> (1989)                                 |
|                        |                                    |                               |                                |           |                    |                 |      | palisade mes. area +23%   |   |
|                        |                                    |                               |                                |           |                    |                 |      | spongy mes. area +39% <sup>†</sup>  |   |
|                        |                                    |                               |                                |           |                    |                 |      | epid. area (x-sect) +25%  |   |
|                        |                                    |                               | 910                            | -         | -                  | -               | -    | leaf thickness +31%   | Vu <i>et al.</i> (1989)                                 |
|                        |                                    |                               |                                |           |                    |                 |      | palisade mes. area +53%   |   |
|                        |                                    |                               |                                |           |                    |                 |      | spongy mes area NS  |   |
|                        |                                    |                               |                                |           |                    |                 |      | epid. area (x-sect) NS  |   |
|                        | GC (45 d)                          | 330                           | 800                            | -         | -                  | -               | -37% | leaf thickness +<br>cell # +<br>intercellular space (-)                     |   |

<sup>s</sup>Plants from all experiments cited in this table were grown in containers, d, days; m, months; y, years.

SLA, specific leaf area (leaf area/total leaf dry weight); GC, growth chamber; GH, glass house; OTC, open top chamber; phy, phytotron; BC, branch chamber.

NS, not significant; LN, low nitrogen; HN, high nitrogen; LF, low fertility; HF, high fertility; <sup>†</sup>, mean of fertility or water treatments.

aT, ambient temperature; XET, xyloglucan endotransglycosylase.

required to produce a given amount of stem tissue in CO<sub>2</sub> enriched atmospheres.

### Leaf expansion

After the primordial leaf is initiated, it grows up into a peg-like leaf axis. This growing leaf axis has an apical meristem at the tip which increases leaf height, and marginal meristems, plate and adaxial meristems which increase leaf width and thickness (Esau 1977). Like all other developing plant organs, growth rates and patterns of leaves are governed by cell division, cell wall loosening, cell wall extensibility, and cell turgor. As mentioned earlier, there has been some discussion concerning which of these processes is most affected by elevated CO<sub>2</sub>. This question is perhaps most pertinent for growth of leaves (as opposed to stems and roots) because they exhibit determinate growth. Any deviation in either cell division or cell expansion could cause significant alterations in final size, anatomy, allometric relationships, and resultant leaf function.

Leaf growth is frequently altered by differences in **plant water potential** (Boyer 1968; Yegappan *et al.* 1982; Taylor *et al.* 1994); water limitations lead to inhibition of both cell expansion and cell division (Jones 1985). This, coupled with the observation that growth in elevated CO<sub>2</sub> enhances the efficiency of water use, has led several authors to attribute increased leaf growth for plants grown in elevated CO<sub>2</sub> to greater cell turgor pressure (Madson 1968). For example, Sasek & Strain (1989) reported that turgor pressure in developing leaves of *Pueraria lobata* was twofold greater in plants grown in elevated CO<sub>2</sub> than for those grown in ambient CO<sub>2</sub> which, they suggested, resulted in increased leaf expansion rates and greater leaf expansivity. However, increased turgor pressure can result in increased cell expansion only when accompanied by cell wall relaxation. In fact, biochemical and molecular properties governing cell wall relaxation and expansivity are thought to be of overriding importance in the control of cell growth (Cosgrove 1993, 1997; Taylor *et al.* 1994).

If greater turgor pressure alone is not sufficient to account for the increases in leaf growth commonly reported for plants grown in elevated CO<sub>2</sub>, then either **cell wall relaxation**, **cell division**, or both must be affected. Examining leaf cell size and number may reveal which of these is most affected by elevated CO<sub>2</sub>. Increased leaf size associated with larger cells suggests that cell expansion has been stimulated, while increased leaf size associated with more cells may imply stimulation of cell division. Indeed, published studies attribute greater leaf growth to increases in cell number (Tables 3, 4 and 5) (*L. corniculatus*, Taylor *et al.* 1994; *Populus* clones,

Ceulemans *et al.* 1995; *Populus*, Gardner *et al.* 1995), and cell size (*Lycopersicum esculentum*, Madson 1968; *Populus* clones, Radoglou & Jarvis 1992; *P. media*, Taylor *et al.* 1994; 3 herbs, Ferris & Taylor 1994) or a combination of both (*Phaseolus vulgaris*, Ranasinghe & Taylor 1996). It appears that no single process has been identified as being responsible for greater leaf lamina size in plants grown in elevated CO<sub>2</sub>. Existing studies, however, do suggest that greater cell expansion may play a larger role in effecting larger leaf size that does enhanced cell division (Murray 1997).

Although studies evaluating numbers and sizes of leaf cells in relation to total leaf size may provide insight into the specific growth processes influenced by CO<sub>2</sub> levels, they do not provide a mechanistic basis for this response. Taylor *et al.* (1994) undertook an explanation of the **cellular mechanisms** driving increased cell expansion in leaves. They examined two species exhibiting different responses to elevated CO<sub>2</sub>; *Plantago media* leaves increase in size due to greater cell size while leaves of *Anthyllis vulneraria* are unresponsive to elevated CO<sub>2</sub>. They found that for *Plantago*, both cell wall plasticity and elasticity were increased in leaves grown in elevated CO<sub>2</sub> implying that cell wall relaxation had occurred. They found no effects of elevated CO<sub>2</sub> on *Anthyllis* cell walls which paralleled its overall lack of response to CO<sub>2</sub> enrichment. Other investigators have temporally separated the processes of cell division and cell expansion in *Phaseolus vulgaris* in an attempt to elucidate the mechanisms underlying increased cell expansion in leaves of plants grown in elevated CO<sub>2</sub> (Taylor *et al.* 1994; Ranasinghe & Taylor 1996). These studies offer three pieces of evidence suggesting that growth in elevated CO<sub>2</sub> caused larger leaf cells by increasing cell wall loosening and extensibility. First, they were able to directly measure increased cell wall extensibility; second, they observed that cell wall yield turgor was reduced; and third, activity of xyloglucan endotransglycosylase (XET) was significantly increased. XET is a putative cell wall loosening enzyme thought to function by cutting and rejoining xyloglucan molecules which connect adjacent microfibrils.

Although data suggest that increases in cell expansion may contribute to larger leaf size more than increased cell division, this is certainly not the case in all species or in the same species at all times of the year (Tables 3, 4 and 5). Ferris *et al.* (1996) reported that in *Lolium perenne*, exposure to elevated CO<sub>2</sub> differentially impacted leaf growth in the spring compared to summer. In the spring, leaf area increased due to increased cell expansion, more epidermal cells per leaf, and increased mesophyll area. However, in the summer, there was a negative effect of elevated CO<sub>2</sub> on leaf cell expansion, epidermal cell length, and mesophyll cell area. There is growing evidence for

**Table 5** Effects of growth in elevated CO<sub>2</sub> on leaf characteristics of native, non-woody species.

| Species                      | §Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf | Leaf<br>area/plant | Leaf<br>#/plant | SLA  | Other leaf<br>Changes  | Reference                   |
|------------------------------|------------------------|-------------------------------|--------------------------------|-----------|--------------------|-----------------|------|--|-----------------------------|
| <i>Desmodium paniculatum</i> | Phy (33 d)             | 350                           | 1000                           | –         | +22%               | –               | –19% | leaf area ratio –36%<br>leaf area duration +34%  | Wulff and Strain (1981)     |
| <i>Lotus corniculatus</i>    | GC (3 m)               | 350                           | 700                            | –         | NS                 | –               | –18% | leaf area ratio –17%<br>leaf weight ratio NS   | Carter <i>et al.</i> (1997) |
|                              | GC (45 d)              | 345                           | 590                            | +37%      | –                  | –               | –    | epid cell # / leaf + sd<br>epid cell # / leaf area +   | Ferris and Taylor (1994)    |
| <i>Sanguisorba minor</i>     | GC (45 d)              | 345                           | 590                            | +31%      | –                  | –               | –    | epid cell # / leaf area +<br>altered leaf shape  |                             |
| <i>Anthyllis vulneraria</i>  | GC (45 d)              | 345                           | 590                            | NS        | –                  | –               | –    | epid cell # / leaf area +  |                             |
| <i>Lonicera japonica</i>     | GC (54 d)              | 350                           | 675                            | –         | +50%               | –               | (–)  | –  | Sasek and Strain (1991)     |
|                              |                        |                               | 1000                           | –         | +50%               | –               | (–)  | –  |                             |
| <i>Lonicera sempervirens</i> | GC (54 d)              | 350                           | 675                            | –         | NS                 | –               | +    | –  |                             |
|                              |                        |                               | 1000                           | –         | NS                 | –               | +    | –  |                             |
| <i>Plantago media</i>        | GC (45 d)              | 345                           | 590                            | +27%      | –                  | –               | –    | epid cell # / leaf area +  | Ferris and Taylor (1994)    |
| <i>Lajia platyglossa</i>     | GC (1 season)          | 300                           | 700                            | –         | –                  | –               | –    | leaf thickness +16%<br>vt. area NS   | St. Omer and Horvath (1984) |
|                              |                        |                               |                                |           |                    |                 |      | phloem area +67%<br>xylem cell diam. NS  |                             |
|                              |                        |                               |                                |           |                    |                 |      | xylem cell # NS  |                             |
|                              |                        |                               |                                |           |                    |                 |      | sieve element diam. +29%   |                             |
| <i>Eichhornia crassipes</i>  | GC (4 wks)             | 330                           | 600                            | NS        | +40%               | +27%            | –    | leaf area index +46%   | Spencer and Bowes (1986)    |
| <i>Taraxacum officinale</i>  | GH (4 m)               | 350                           | 700                            | NS        | NS                 | NS              | –64% | altered areaperimeter<br>altered length:width  | Thomas and Bazzaz (1996)    |
|                              |                        |                               |                                |           |                    |                 |      | leaves "boothier"  |                             |
| <i>Pueraria lobata</i>       | Phy (45 d)             | 350                           | 675                            | +         | +22%               | –               | –    | If. expansion rt. +40%   | Sasek and Strain (1989)     |
|                              |                        |                               | 1000                           | +         | +54%               | –               | –    | If. production rt. +12%<br>(above values are averaged for<br>the two CO <sub>2</sub> levels) |                             |
| <i>Abutilon theophrasti</i>  | GH (53 d)              | 360                           | 700                            | –         | NS                 | –               | NS   | vertical distribution of<br>leaf area NS   | Hirose <i>et al.</i> (1996) |

| Species                        | <sup>§</sup> Location/<br>Duration | Ambient<br>[CO <sub>2</sub> ] | Elevated<br>[CO <sub>2</sub> ] | Area/leaf | Leaf<br>area/plant                 | Leaf<br>#/plant   | SLA | Other leaf<br>Changes                                       | Reference                     |
|--------------------------------|------------------------------------|-------------------------------|--------------------------------|-----------|------------------------------------|-------------------|-----|---|-------------------------------|
| <i>Ambrosia artemisiifolia</i> | GH (1 season)                      | 350                           | 700                            | -         | (-)                                | -                 | (-) | leaf area ratio +<br>plastochron index NS                   | Carbutt <i>et al.</i> (1990)  |
|                                | GH (31 d)                          | 400                           | 700                            | -         | 18°C: NS<br>28°C: +55%<br>38°C: NS | -                 | -   |   | Ackerly <i>et al.</i> (1992)  |
|                                | GH (53 d)                          | 360                           | 700                            | -         | NS                                 | -                 | NS  | vertical distribution of<br>leaf area NS                    | Hirose <i>et al.</i> (1996)   |
| <i>Agrostis capillaris</i>     | GH (1 season)                      | 350                           | 700                            | -         | (-)                                | -                 | (-) | leaf area ratio +<br>growth rate increased                  | Carbutt <i>et al.</i> (1990)  |
|                                | GH (23 wks)                        | 360                           | 610                            | (-)       | NS                                 | +39%*             | -   | then decreased  | Newberry and Wolfenden (1996) |
| <i>Calamagrostis epigetos</i>  | GC (1 m)                           | 350                           | 700                            | -         | -                                  | -                 | (-) | leaf length +<br>tiller # +36%<br>leaf area ratio (-)       | Gloser and Barták (1994)      |
| <i>Nymphaea odorata</i>        | OTC (5 m)                          | 350                           | 650                            | +18%      | -                                  | +75% <sup>¶</sup> | -   | relative girth rate +32%<br>leaf number × lifespan<br>+116% | Idso <i>et al.</i> (1990)     |

<sup>§</sup>Plants from all experiments cited in this table were grown in containers. d, days; m, months; y, years.

SLA, specific leaf area (leaf area/total leaf dry weight); GC, growth chamber; GH, glass house; OTC, open top chamber; phy, phytotron; BC, branch chamber.

NA, not significant; LN, low nitrogen; HN, high nitrogen; LF, low fertility; HF, high fertility; ¶, not statistically significant; \*, mean of fertility or water treatments

some species that elevated CO<sub>2</sub> stimulates early expansion of leaves but that the effect then diminishes with time ultimately resulting in leaves of similar size (*Pinus taeda*, Tolley & Strain 1984; *Populus*, Radoglou & Jarvis 1990a; *Populus* clones Taylor *et al.* 1994; *Pinus palustris*, Pritchard *et al.* 1998; *Glycine max*, Sims *et al.* 1998). Nevertheless, increased rates of leaf expansion could significantly impact total plant productivity even if leaves do eventually reach identical fully expanded size (Sasek & Strain 1989).

#### Lamina size

Regardless of the cellular mechanisms involved, exposure to elevated CO<sub>2</sub> more often than not results in increased **area per leaf**, greater **leaf thickness**, more **leaves per plant**, and higher **total leaf area per plant** (Tables 3, 4 and 5). In 19 reports of surface area per leaf, 58% exhibited greater area per leaf, 37% were not affected, and 11% had decreased leaf area. In two other reports increased area per leaf was reported in fertile soil but was not observed when plants were not fertilized. In 16 reports in which leaf thickness was measured, thickness usually increased (81%), but sometimes was unaffected (19%) by elevated CO<sub>2</sub>. For 63 observations of total leaf area per plant, growth in elevated CO<sub>2</sub> resulted in increases more than half the time (57%), sometimes resulted in no change (25%), and 10% decreased; 5% exhibited increases when fertilized and no effect when unfertilized, and 3% of species exhibited increased leaf area at 28 °C but decreased area at 38 °C. Leaf area increased for one species during the spring, but decreased during the summer. The average net increase in leaf area per plant was 24%. Crop species exhibited the greatest average increase in whole plant leaf area (+37%) compared to trees (+14%) and wild, nonwoody plants (+15%). In addition to altered LAR, LWR, leaf area per plant and leaf size, decreases in specific leaf area (SLA=leaf area/total leaf dry weight) have often been the result of altered anatomy or increased starch accumulation. Of 49 observations, 78% reported a decrease in SLA, 18% showed no significant difference, and 4% increased. This resulted in an average net decrease in SLA (−16%). Tree species and wild, nonwoody species exhibited the most appreciable reduction in SLA (−14% and −20%; Tables 1, 2) compared to crop species (−6%; Table 3). It is generally thought that decreases in specific leaf area result from increased accumulation of leaf total nonstructural carbohydrates (TNC), and accumulation of TNC occurs when C fixation exceeds C utilization. Therefore, crop species would be expected to accumulate less TNC because of rapid growth rates and high sink activity than more slowly growing, nutrient limited natural species.

#### Leaf ultrastructure

Along with biochemical and biophysical studies, ultrastructural data may provide useful information about cellular processes that drive increased leaf growth at high levels of CO<sub>2</sub>. Robertson & Leech (1995) reported increased leaf growth rate in 7-day-old *Triticum aestivum* seedlings which they attributed to increased cell and chloroplast expansion. Although cell and chloroplast profile (cross-sectional) areas were greater, the number, positioning, shape and internal organization (granal stacking) of chloroplasts were unaffected by elevated CO<sub>2</sub>. However, chloroplasts from leaves developing in elevated CO<sub>2</sub> contained far less starch than those grown in ambient conditions. An increase in mitochondrial biogenesis in basal leaf cells after only 12 h postmitosis (Robertson *et al.* 1995) suggested that this decrease in starch was probably due to increased rates of respiration. The decrease in chloroplast starch at 7 days resulting from growth in elevated CO<sub>2</sub> was not observed in older (4 weeks) wheat plants. Studies of mature leaves exposed to elevated CO<sub>2</sub> often report more and larger starch grains which in some cases may inhibit **chloroplast structure and function**. For example, several studies have attributed the phenomenon of **photosynthetic acclimation** to elevated CO<sub>2</sub> to disruptions in chloroplast integrity caused by excessive starch accumulation (Cave *et al.* 1981; Wulff & Strain 1981; Ehret & Jolliffe 1985; Yelle *et al.* 1989). Reductions in granal stacking and/or mechanical disruption of chloroplast structure induced by elevated CO<sub>2</sub> may, in some species, only occur when soil N is limiting (Kutik *et al.* 1995), or when both soil N and water are limiting (Pritchard *et al.* 1997). Investigating leaf cell ultrastructure over time, beginning with leaf initiation and ending with leaf senescence, might prove useful in elucidating the cellular mechanisms involved in leaf expansion, and further, could provide insight into photosynthetic acclimation and the general decrease in response to elevated CO<sub>2</sub> reported to occur over time.

#### Leaf anatomy

In addition to altered leaf expansion (resulting in changes in leaf size, number, total leaf area per plant, and ultrastructure), changes in internal anatomy and leaf shape are often observed in plants grown in elevated CO<sub>2</sub> (Tables 3, 4, and 5). For example, an extra layer of palisade cells has been observed in *Glycine max* (Rogers, Thomas & Bingham 1983; Thomas & Harvey 1983; Vu *et al.* 1989) and *Castanea sativa* (Mousseau & Enoch 1989) grown in elevated CO<sub>2</sub>. Studies typically report increased total **mesophyll cross-sectional area** in leaves from plants grown in elevated CO<sub>2</sub> (*Pinus radiata*, Conroy *et al.* 1986; *Populus trichocarpa*, Radoglou & Jarvis 1990a; *Populus*,

Radoglou & Jarvis 1992) although reductions have also been reported (*Pinus ponderosa*, Pushnik *et al.* 1995). However, effects of elevated CO<sub>2</sub> on leaf anatomy may vary depending on stage of leaf development, soil fertility, and season of the year. For instance, Pritchard *et al.* (1998) found that in the early stages of needle development in *Pinus palustris*, elevated CO<sub>2</sub> increased needle fascicle volume by 8% and cross-sectional area by 15% due to increased transfusion tissue area (3%), mesophyll area (17%), and vascular tissue area (7%). But in later stages, fascicle volume was reduced 8% resulting from 7% shorter needles and reduced mesophyll (19%), vascular tissue (10%), and epidermal (19%) cross-sectional areas. Conroy *et al.* (1986) reported that for *Pinus radiata*, mesophyll area was increased by elevated CO<sub>2</sub> when P was nonlimiting, but was not affected when P was limiting. Ferris *et al.* (1996) studied *Lolium perenne*; mesophyll area was increased by high CO<sub>2</sub> concentration in spring, but was decreased in summer.

**Vascular tissue area** has also been reported to increase in leaves (*Pinus taeda*, Thomas & Harvey 1983; *Pinus radiata*, Conroy *et al.* 1986; *Pinus ponderosa*, Pushnik *et al.* 1995; *Glycine max*, Leadley *et al.* 1987; *Layia platyglossa*, St. Omer & Horvath 1984) and leaf petioles (*Lypersicon esculentum*, Ho 1977). However, Pritchard *et al.* (1997) observed a trend for reduced phloem area in needles of *Pinus palustris* due to fewer, not smaller, cells. They explained these atypical results by suggesting that the production of secondary phloem in the two-year-old needles sampled was negatively impacted by growth in elevated CO<sub>2</sub>. The other studies on pine were conducted for shorter durations. Clearly, the lack of data prohibits generalizations about these patterns in either crops or other species.

Increased mesophyll and vascular tissue area commonly reported may be important determinates of both photosynthetic and assimilate transport capacity. However, examining **allometric relationships** between tissue types may be of more use in evaluating the effects of elevated CO<sub>2</sub> on leaf function than studies which simply measure leaf thickness. In one study of these relationships, Leadley *et al.* (1987) reported that although *Glycine max* leaves were thicker when grown in elevated CO<sub>2</sub>, they had less palisade cell surface area per unit of leaf area. Internal cell surface area exposed to intercellular spaces is highly correlated with photosynthesis and water use. Additionally, they found that there was a greater ratio of vascular tissue to total leaf cross-sectional area in leaves from plants growing in elevated CO<sub>2</sub>. Pushnik *et al.* (1995) also observed an increase in the ratio of vascular tissue cross-sectional area to total needle cross-sectional area in *Pinus ponderosa*. Pritchard *et al.* (1998) reported that for *Pinus palustris*, CO<sub>2</sub> enhancement resulted in a 17% reduction in mesophyll cell surface area

per unit of needle volume when N was limiting. No effects, however, were observed on the proportion of needle volume allocated to a given tissue type at any needle age. In other studies in which anatomical allometry was examined, Radoglou & Jarvis (1990a) observed no effect of increased CO<sub>2</sub> level on the ratios of palisade to spongy parenchyma, spongy parenchyma to total leaf thickness, or palisade parenchyma to total leaf thickness in *Populus*.

### Stomates

Many studies have reported that growth in elevated CO<sub>2</sub> alters stomatal characteristics. Stomatal density has been observed to increase (Thomas & Harvey 1983; Gaudillière & Mousseau 1989), decrease (Woodward & Bazzaz 1988), or stay the same (Mousseau & Enoch 1989; Radoglou & Jarvis 1990b, 1992; Estiarte *et al.* 1994; Pritchard *et al.* 1998) in plants grown in elevated atmospheric CO<sub>2</sub>. Beerling & Woodward (1995) looked at results from historical studies (preindustrial herbarium specimens were compared to contemporary leaves) and experimental studies, and reported that 60% of species show a decline in stomatal density due to elevated CO<sub>2</sub>; this number increased to 85% when only experimental data were considered. In addition to altered stomatal frequencies, stomatal patterning may be modified. Boetsch *et al.* (1996) reported that extra subsidiary cells were associated with nearly half of stomatal complexes in *Tradescantia* leaves grown in elevated CO<sub>2</sub> which suggests that recruitment of epidermal cells into stomatal complexes was stimulated. However, Murray (1995) hypothesized that changes in stomatal conductance and WUE resulting from CO<sub>2</sub> enrichment are probably the result of adjustments in stomatal apertures, not consistent morphogenic effects on stomatal abundance. Examination of guard/subsidiary cell ultrastructure, anatomy, and ontogeny with simultaneous observations of gas exchange may provide a more complete understanding of CO<sub>2</sub> effects on biophysical and biochemical aspects of stomatal functioning. Modelling future responses of vegetation will require a sound understanding of how stomatal anatomy and function are affected by higher CO<sub>2</sub> levels (Wagner *et al.* 1996).

### Leaf shape

Few studies have examined the effects of elevated CO<sub>2</sub> on leaf shape even though morphological changes in leaf shape may be of greater functional significance than changes in leaf level photosynthesis (Niklas & Owens 1989; Niklas 1989; Thomas & Bazzaz 1996). There is some evidence that carbohydrate availability may be critical in determining leaf form, at least in species exhibiting

heteroblastic leaf development (Thomas & Bazzaz 1996). Moreover, Ranasinghe & Taylor (1996) found that spatial patterns of cell wall extensibility were altered in leaves from plants grown in elevated CO<sub>2</sub> which they suggested could lead to alterations in leaf shape. Thomas & Bazzaz (1996) observed shifts in allometric relationships between leaf area and perimeter, and between leaf length and width in *Taraxicum officinale*; and between leaf width and area, leaf length and area, and leaf length and width in *Plantago major* exposed to elevated CO<sub>2</sub> which caused leaves to be more dissected. In contrast, Leadley & Reynolds (1989) found no differences in allometric relationships among length, width, and area of *Glycine max* leaves. It is clear that more attention should be paid to leaf shape in studies examining effects of elevated CO<sub>2</sub> on plant structure and function.

**In conclusion**, although growth in CO<sub>2</sub>-enriched atmospheres results in more leaves per plant, leaf initiation is usually reduced relative to whole plant growth. Although highly variable, rates and magnitude of leaf expansion are enhanced (at least temporarily); this results more often from increased cell expansion than increased cell division. Increased expansion appears to result from greater cell wall relaxation and/or greater cell turgor. Stomatal densities usually decrease, and mesophyll and vascular tissue cross-sectional areas increase contributing to greater total leaf thickness. Although thicker, leaf organization may suggest structural shifts that may limit plant capacity to assimilate carbon (acclimation). Very little is known concerning the impact of elevated CO<sub>2</sub> on leaf shape, or if growth in elevated CO<sub>2</sub> will differentially influence different leaf types (for example, compound vs. simple, opposite vs. alternate).

## Root development

### *Primary growth of roots*

Plant roots play a crucial role as the interface between lithosphere and biosphere. Spatial and temporal root structural characteristics govern both plant and soil processes including: (i) root weathering of soil; (ii) input of carbon to soil; (iii) mining soil for resources; and (iv) erosion (Rogers *et al.* 1992). And although roots often exhibit the greatest relative increase in biomass of all plant organs when grown in elevated CO<sub>2</sub>, there are few studies on root structural responses, and thus these responses are poorly characterized (Rogers *et al.* 1994; 1997b; Rogers *et al.* 1997a). Reviews have evaluated the effects of elevated CO<sub>2</sub> on roots, and each echoes the need to further investigate below-ground processes, including root structure (Stulen & den Hertog 1993; Rogers *et al.* 1994; Rogers *et al.* 1997b). For a comprehen-

sive tabular summary of CO<sub>2</sub> effects on roots please refer to Rogers *et al.* 1994.

The **root apical meristem** is structurally and functionally different from the shoot apical meristem. In shoots, branching patterns are determined by events at the shoot apical meristem, whereas in roots, lateral organs are initiated in the pericycle, some distance distal to the root tip (Taylor 1997). Functionally, the root axis exhibits negative gravitropism while the shoot axis is positively gravitropic. Finally, roots are thought to play a major role in regulating the growth of above-ground plant organs. Roots sense the availability of soil water and nutrients, and accordingly alter production and transport of hormones such as cytokinins and ABA to shoots, thereby modulating the activity of meristematic tissues above the ground, as well as expression of genes coding for photosynthetic enzymes phosphoenolpyruvate carboxylase, carbonic anhydrase, and the small Rubisco subunit (Aiken & Smucker 1996).

Because of structural and functional differences between the shoot apical meristem and the root apical meristem, one might expect that the effects of elevated CO<sub>2</sub> on components of development leading to increased shoot growth and altered patterns of branching above-ground may not be analogous to below-ground processes. Ferris & Taylor (1994) suggested that cell expansion may be stimulated to a greater extent than cell division in roots. They observed increased root extension rates in *Sanguisorba minor*, *Lotus corniculatus*, *Anthyllis vulneraria*, and *Plantago media* grown in elevated CO<sub>2</sub>. They found that cell length was unaffected by CO<sub>2</sub> treatment from  $\approx 15\text{--}25\mu\text{m}$  up to  $\approx 1\text{mm}$  behind the growing tip. Beyond 1 mm, cell length increased at a greater rate in all four species grown in elevated compared to ambient CO<sub>2</sub>. Also, they observed increased turgor pressure in plants grown in elevated compared to ambient CO<sub>2</sub>. They concluded that stimulation of root growth was the result of increased cell expansion caused by cell wall loosening, in concert with higher cell turgor pressure, rather than by increased cell division. Crookshanks *et al.* (1998) reported a 40% greater mean root cortical cell length in elevated compared to ambient CO<sub>2</sub> at a distance of 375  $\mu\text{m}$  from the root tip which they attributed to increased cell wall extensibility but final cortical cell length, however, was unaltered by exposure to elevated CO<sub>2</sub>. Kinsman *et al.* (1997) reported that increased root growth in *Dactylis glomerata* was mainly the result of a greater proportion of dividing cells in the apical meristem. Although they also observed greater mitotic indices indicative of shortened cell cycles, these effects were not of sufficient magnitude to cause significant changes in root growth. As shown by the contra-



dictory results of these three studies, further work is needed to elucidate the cellular mechanisms underlying effects of elevated atmospheric CO<sub>2</sub> on stimulation of root elongation behind the apical meristem and also on the events controlling stimulation of lateral root formation in the pericycle. Use of *Arabidopsis thaliana* mutants to elucidate the specific cellular events leading to shifts in root growth as described by Crookshanks *et al.* (1998) currently holds great promise [see also Schiefelbein *et al.* (1997) concerning use of *Arabidopsis* mutants to study mechanistic controls on root development].

Regardless of the cellular mechanisms involved, exposure of the plant canopy to elevated CO<sub>2</sub> usually stimulates growth of roots. Rogers *et al.* (1992) reported a 27% increase in **root diameter** in the root hair zone, a 23% increase in stele diameter, and a 28% increase in cortex width in *Glycine max* grown in elevated CO<sub>2</sub>. Increased root diameters have also been reported for *Pinus taeda* (Larigauderie *et al.* 1994). These results notwithstanding, St. Omer & Horvath (1984) found no difference in stele diameter, diameter of tracheary elements, or wall thickness of tracheary elements of *Layia platyglossa* roots grown at higher than ambient CO<sub>2</sub> levels.

Besides greater root diameters, increased **total root lengths** are often observed in plants grown in elevated CO<sub>2</sub> (*Glycine max*, Rogers *et al.* 1992; *Senecio vulgaris*, Bernston & Woodward 1992; *Trifolium repens*, Jongen, Fay & Jones 1996). This may not be true for all species or for roots at all depths within the same species. For example, in *Pinus taeda* grown in elevated CO<sub>2</sub>, the upper lateral root fraction increased but the proportion of remaining root components generally declined (Larigauderie *et al.* 1994). Mo *et al.* (1992) reported that, although there was no effect of elevated CO<sub>2</sub> on dry weight in the 0–40 cm depth range, root length in the 0–10 cm range decreased 31% in an assemblage of tallgrass prairie species.

Differential effects of CO<sub>2</sub> concentration on root branching may lead to altered root architecture and altered ability of roots to acquire water and nutrients from the soil profile. For example, Bernston & Woodward (1992) observed for *Senecio vulgaris* that exposure to elevated CO<sub>2</sub> caused more horizontal branching angles of roots contributing to greater horizontal root spread, and Rogers *et al.* (1992) observed longer second-order laterals in *Glycine max*, which could lead to deeper root penetration and thus greater exploration of soil for nutrients and water. Conversely, Del Castillo *et al.* (1989) reported that *Glycine max* grown in elevated CO<sub>2</sub> had more roots due to increased branching rather than longer roots. Increased root numbers may enable plants to more

efficiently explore and mine the same volume of soil instead of stimulating exploration into soil deeper or further away.

It is important that more studies focus on effects of elevated CO<sub>2</sub> on root system architecture. The extent of root branching has major implications for the efficiency of water and mineral extraction from soil. Additionally, altered rooting patterns may contribute to root overlap between adjacent plants, possibly intensifying below-ground competition (Bernston & Woodward 1992). Perhaps the 31% reduction in root length in the top 10 cm of soil in the assemblage of prairie species resulting from elevated CO<sub>2</sub> (Mo *et al.* 1992) portends altered root competition.

## Limitations on carbon assimilation

### *Limitations imposed by physiology or structure*

This review has focused primarily on the ways in which increased plant carbon assimilation may alter developmental processes and ultimately plant structure. Different plant species, however, are not able to assimilate carbon to the same extent. Limitations may result from differing physiological strategies employed to fix carbon (e.g. C3 vs. C4 vs. CAM photosynthesis) (Poorter 1993), or from differential capacity to metabolize carbohydrates by actively growing plant parts. For example, plants which have rapid growth rates or very large storage organs are not as prone to photosynthetic acclimation to elevated CO<sub>2</sub> as slow-growing species with weaker sinks. Photosynthesis, as with most other biochemical processes in living organisms, is subject to end-product inhibition. If plants are unable to utilize fixed carbon for growth, the result is typically an increase in total nonstructural carbohydrates (and perhaps increased C-based secondary compounds) (Poorter *et al.* 1997). Leaf starch accumulation, as discussed earlier, may inhibit photosynthesis by mechanically altering the integrity of chloroplasts, and soluble sugars may inhibit photosynthesis by biochemical feedback processes acting at the level of gene expression. Feedback inhibition of photosynthesis may include adjustments in Rubisco amount or activity, reductions in components of photosystem II (Pennanen *et al.* 1993; Van Oosten *et al.* 1994), reductions in thylakoid stacking within chloroplasts (Wulff & Strain 1981; Kutík *et al.* 1995; Pritchard *et al.* 1997), or a shortage of cytosolic inorganic phosphates (Stitt 1991).

In addition to limitations on carbon assimilation imposed by mechanical and biochemical processes within source leaves and from limitations resulting from weak sink activity, there may be other **structural/**

**functional attributes that may also limit plant capacity to exploit the extra carbon** available in a higher CO<sub>2</sub> world. For example, Körner *et al.* (1995) found that, following exposure to elevated CO<sub>2</sub>, species which load phloem symplastically accumulated 41% total leaf non-structural carbohydrates compared to 25% in species which exhibit apoplastic phloem loading (see also Poorter *et al.* 1997). This implies that plant species may be 'predisposed' to respond to increasing atmospheric CO<sub>2</sub> based not only on photosynthetic capacity and sink strength, but also on the efficiency of assimilate transport from sources to sinks. Pritchard *et al.* (1997) suggested that differences in structure (e.g. vein configurations, plasmodesmatal connectivity, sieve cell organization and form, absence or presence of P-protein), that reflect different strategies for short-distance assimilate transport, phloem loading, and long-distance phloem transport, may, in part, account for observed differences in response patterns of woody broadleaf plants and conifers (Ceulemans & Mousseau 1994; Pushnik *et al.* 1995). Clearly, future studies should attempt to understand plant response to elevated CO<sub>2</sub> not only in the context of plant physiological strategy (photosynthetic pathway) or patterns of biomass accumulation, but also in the context of plant structural attributes.

#### *Limitations arising as experimental artifacts*

In almost all of the studies reviewed here, plants were grown individually in pots. This could cause concern for two reasons: (i) first, container grown plants may exhibit greater down-regulation in rates of photosynthesis due to a source-sink imbalance resulting from root constriction (Arp 1991), and/or from nutrient limitations that sometimes plague pot studies (McConnaughay *et al.* 1993), or (ii) individually grown plants may exhibit greater plasticity than plants grown in competition. Furthermore, the tree species studied were typically seedlings and therefore, may not accurately reflect structural responses of mature trees. Moreover, in many cases, plant growth responses to elevated CO<sub>2</sub> may be mediated by nutrient availability (Conroy *et al.* 1990b; Prior *et al.* 1997; Sims *et al.* 1998) which many published studies have not adequately addressed. It is becoming evident that experimental approaches must go beyond simply evaluating the response of isolated container grown plants to elevated CO<sub>2</sub> levels in favour of in-ground studies in which plants are grown at densities reflective of either the natural or agricultural ecosystems where they typically occur. Recently, FACE (free air CO<sub>2</sub> enrichment) studies, and open top field chambers are beginning to be used more regularly to conduct multi-factor in-ground experiments or to study entire ecosystems (see Saxe *et al.* 1998 for a recent discussion on

experimental methodology). These studies will fill important gaps in our understanding of plant response to global change, and these new data will likely modify our existing understanding of plant physiological and structural responses to elevated CO<sub>2</sub> levels.

#### **Research recommendations**

Review of the extant data on the influence of elevated atmospheric CO<sub>2</sub> levels on plants has revealed the need for focused efforts on nearly every aspect of plant development and structure. We recommend that the following be taken into consideration in future research on the effects of elevated CO<sub>2</sub> on plant structure.

- 1 How shifts in cellular (cell division and cell expansion) and higher level (morphogenesis and histogenesis) growth processes contribute to alterations in leaf, branch, stem, and root structure.
- 2 Whether increased cell division is driven by greater rates of cell expansion or is transduced via molecular cues such as sucrose, cyclins, XET, expansins, cytokinins or other molecules important in controlling the cell cycle and cell wall mechanical properties.
- 3 How cell patterning is altered during morphogenesis and histogenesis resulting in plant organs with different shapes and anatomical organizations, and how cells are partitioned to branch and leaf primordia vs. internodes.
- 4 Determine if whole plant growth responses to elevated CO<sub>2</sub> may be predictable based on specific developmental events occurring within meristematic tissues. This may prove useful in the search for plant functional types.
- 5 Alterations in plant allometry including shifts in scaling relationships evident at subcellular (e.g. chloroplasts:mitochondria), anatomical, and morphological levels. Structural alterations in plant organs resulting from growth in elevated CO<sub>2</sub> must be considered in the context of whole plant responses instead of considering single structures or processes in isolation.
- 6 Anatomical studies of primary and secondary stem and branch anatomy in conjunction with measures of hydraulic conductivity will facilitate a more complete understanding of plant resistance to drought and frost as well as revealing potential shifts in functional relationships of leaves and stems.
- 7 It has been suggested that compound leaves have a much greater capacity for indeterminate growth than simple leaves (Poethig 1997). Is there a differential CO<sub>2</sub> response between plants with compound vs. simple leaves?
- 8 The role of ultrastructural, anatomical, and morphological leaf adjustments in the downregulation of photosynthesis rates often observed after prolonged exposure to elevated CO<sub>2</sub>.

9 Alterations in plant micromorphology including, but not limited to, root hair and leaf trichome density and structure. Trichomes are vital in protecting plants against pathogens and herbivores and root hair characteristics have a major influence on water and mineral uptake.

## Conclusions

It is clear from this review that the effects of elevated CO<sub>2</sub> on plant development and structure are both many and varied. The necessity of understanding the influence of growth in elevated CO<sub>2</sub> on cellular developmental processes, and bridging these basic growth mechanisms to higher level structure and function, is emerging. Ackerly *et al.* (1992) were correct in their assertion that 'elucidation of the relationship between individual developmental processes and whole plant growth has proven much more difficult than the comparable analysis of the mechanistic basis of carbon assimilation and water relations.' However, in spite of these difficulties, several recent studies have provided inroads towards disentangling the effects of elevated CO<sub>2</sub> on the separate but interdependent processes governing growth and development including cell division, cell expansion, primordium initiation, cell differentiation, and organogenesis. The most significant direct effect of elevated CO<sub>2</sub> on plant growth is certainly an increase in carbohydrate availability and increased water-use efficiency. Ultimately, both increased carbon and more efficient water use combine to stimulate cell proliferation either by promoting cell division, cell expansion, or both. Atmospheric CO<sub>2</sub> levels predicted for the next century therefore will likely result in faster seasonal, and successional canopy development and closure. The ability of a given species within the canopy to exploit extra carbon, however, will largely be a function of its inherent physiological and structural attributes integrated with anatomical/morphological plasticity. Some species are likely to overtop others. A more thorough mechanistic understanding of the ways elevated CO<sub>2</sub> will impact structure will emerge as plant biologists develop a better knowledge of how plant developmental processes are regulated.

## Acknowledgements

This work is supported by Interagency Agreement No. DE-AI05-95ER62088 from the U.S. Department of Energy, Environmental Sciences Division and by the Experimental Program to Stimulate Competitive Research, U.S. Environmental Protection Agency, Contract No. R826259-01. Any opinions, findings, and conclusions or recommendations expressed in this publication are those of the authors and do not necessarily reflect the views of the U.S. Department of Energy or the U.S. Environmental Protection Agency. The

authors thank R.R. Dute, G.B. Runion, and M.L. Davis for critical reviews of this manuscript.

## References

- Ackerly DD, Coleman JS, Morse SR, Bazzaz FA (1992) CO<sub>2</sub> and temperature effects on leaf area production in two annual plant species. *Ecology*, **73**, 1260–1269.
- Aiken RM, Smucker AJM (1996) Root system regulation of whole plant growth. *Annual Review of Phytopathology*, **34**, 325–346.
- Allen LH Jr, Amthor JS (1995) Plant physiological responses to elevated CO<sub>2</sub>, temperature, air pollution, and UV-B radiation. In: *Biotic Feedbacks in the Global Climatic System: Will the Warming Feed the Warming?* (eds Woodwell GM, Mackenzie FT), pp. 51–84. Oxford University Press, New York.
- Amthor JS (1995) Terrestrial higher-plant response to increasing atmospheric [CO<sub>2</sub>] in relation to the global carbon cycle. *Global Change Biology*, **1**, 243–274.
- Anderson AS (1976) Regulation of apical dominance by ethephon, irradiance and CO<sub>2</sub>. *Physiologia Plantarum*, **37**, 303–308.
- Arp WJ (1991) Effects of source-sink relations on photosynthetic acclimation to elevated CO<sub>2</sub>. *Plant, Cell and Environment*, **14**, 869–875.
- Ashton PMS, Berlyn GP (1994) A comparison of leaf physiology and anatomy of *Quercus* environments. *American Journal of Botany*, **81**, 589–597.
- Atkinson CJ, Taylor JM (1996) Effects of elevated CO<sub>2</sub> on stem growth, vessel area and hydraulic conductivity of oak and cherry seedlings. *New Phytologist*, **133**, 617–626.
- Atkinson CJ, Taylor JM, Wilkins D, Besford RT (1997) Effects of elevated CO<sub>2</sub> on chloroplast components, gas exchange and growth of oak and cherry. *Tree Physiology*, **17**, 319–325.
- Bazzaz FA (1990) The response of natural ecosystems to the rising global CO<sub>2</sub> levels. *Annual Review of Ecology and Systematics*, **21**, 167–196.
- Beerling DJ, Woodward FI (1995) Stomatal responses of variegated leaves to CO<sub>2</sub> enrichment. *Annals of Botany*, **75**, 507–511.
- Bernston GM, Woodward FI (1992) The root system architecture and development of *Senecio vulgaris* in elevated CO<sub>2</sub> and drought. *Functional Ecology*, **6**, 324–333.
- Bezemer TM, Jones TH (1998) Plant–insect herbivore interactions in elevated atmospheric CO<sub>2</sub>: quantitative analysis and guild effects. *Oikos*, **82**, 212–222.
- Boetsch J, Chin J, Ling M, Croxdale J (1996) Elevated carbon dioxide affects the patterning of subsidiary cells in *Tradescantia* stomatal complexes. *Journal of Experimental Botany*, **47**, 935–931.
- Bowes G (1991) Growth at elevated CO<sub>2</sub>: photosynthetic responses mediated through Rubisco. *Plant, Cell and Environment*, **14**, 795–806.
- Boyer JS (1968) Relationship of water potential to growth of leaves. *Plant Physiology*, **43**, 1056–1062.
- Brown K, Higginbotham KO (1986) Effects of carbon dioxide enrichment and nitrogen supply on growth of boreal tree seedlings. *Tree Physiology*, **2**, 223–232.
- Carter EB, Theodorou MK, Morris P (1997) Responses of *Lotus corniculatus* to environmental change. *New Phytologist*, **136**, 245–253.

- Cave G, Tolley LC, Strain BR (1981) Effect of carbon dioxide enrichment on chlorophyll content, starch content and starch grain structure in *Trifolium subterraneum* leaves. *Physiologia Plantarum*, **51**, 171–174.
- Ceulemans R, Mousseau M (1994) Effects of elevated atmospheric CO<sub>2</sub> on woody plants. *New Phytologist*, **127**, 425–446.
- Ceulemans R, Van Praet L, Jiang XN (1995) Effects of elevated CO<sub>2</sub> enrichment, leaf position and clone on stomatal index and epidermal cell density in poplar (*Populus*). *New Phytologist*, **131**, 99–107.
- Chapin FS III, Bloom AJ, Field CB, Waring RH (1987) Plant responses to multiple environmental factors. *Biosciences*, **41**, 29–36.
- Cheeseman JM, Barreiro R, Lexa M (1996) Plant growth modelling and the integration of shoot and root activities without communicating messengers: Opinion. *Plant and Soil*, **185**, 51–64.
- Clark SE (1997) Organ formation at the vegetative shoot meristem. *The Plant Cell*, **9**, 1067–1076.
- Conroy J, Barlow EWR, Bevege DI (1986) Response of *Pinus radiata* seedlings to carbon dioxide enrichment at different levels of water and phosphorus: growth, morphology and anatomy. *Annals of Botany*, **57**, 165–177.
- Conroy JP, Milham PJ, Mazur M, Barlow EW (1990a) Growth, dry weight partitioning and wood properties of *Pinus radiata* D. Don after 2 years of CO<sub>2</sub> enrichment. *Plant, Cell and Environment*, **13**, 329–337.
- Conroy JP, Milham PJ, Reed ML, Barlow EW (1990b) Increases in phosphorus requirements for CO<sub>2</sub>-enriched pine species. *Plant Physiology*, **92**, 977–982.
- Cosgrove DJ (1993) Tansley Review no. 46. Wall extensibility: its nature, measurement and relationship to plant cell growth. *New Phytologist*, **124**, 1–23.
- Cosgrove DJ (1997) Relaxation in a high-stress environment: The molecular basis of extensible cell walls and cell enlargement. *The Plant Cell*, **9**, 1031–1041.
- Crookshanks M, Taylor G, Dolan L (1998) A model system to study the effects of elevated CO<sub>2</sub> on the developmental physiology of roots: the use of *Arabidopsis thaliana*. *Journal of Experimental Botany*, **49**, 593–597.
- Curtis PS, Vogel CS, Pregitzer KS, Zak DR, Teeri JA (1995) Interacting effects of soil fertility and atmospheric CO<sub>2</sub> on leaf area growth and carbon gain physiology in *Populus x euramericana* (Dode) Guinier. *New Phytologist*, **129**, 253–263.
- Del Castillo D, Acock B, Reddy VR, Acock MC (1989) Elongation and branching of roots on soybean plants in a carbon dioxide-enriched aerial environment. *Agronomy Journal*, **81**, 692–695.
- Díaz S (1995) Elevated CO<sub>2</sub> responsiveness, interactions at the community level and plant functional types. *Journal of Biogeography*, **22**, 289–295.
- Donaldson LA, Hollinger D, Middleton TM, Souter ED (1987) Effect of CO<sub>2</sub> enrichment on wood structure in *Pinus radiata* D. Don. *IAWA Bulletin*, **8**, 285–289.
- Downton WJS, Grant WJR, Chacko EK (1990) Effect of elevated carbon dioxide on the photosynthesis and early growth of mangosteen (*Garcinia mangostana* L.). *Scientia Horticulturae*, **44**, 215–225.
- Ehret DL, Jolliffe PA (1985) Leaf injury to bean plants grown in carbon dioxide enriched atmospheres. *Canadian Journal of Botany*, **63**, 2015–2020.
- El Kohen A, Mousseau M (1994) Interactive effects of elevated CO<sub>2</sub> and mineral nutrition on growth and CO<sub>2</sub> exchange of sweet chestnut seedlings (*Castanea sativa*). *Tree Physiology*, **14**, 679–690.
- El Kohen A, Rouhier H, Mousseau M (1992) Changes in dry weight and nitrogen partitioning induced by elevated CO<sub>2</sub> depend on soil nutrient availability in sweet chestnut (*Castanea sativa* Mill). *Annales Des Sciences Forestières*, **49**, 1–8.
- Esau K (1977) *Anatomy of Seed Plants*, 2nd edn. John Wiley, Chichester.
- Estiarte M, Peñuelas J, Kimball BA *et al.* (1994) Elevated CO<sub>2</sub> effects on stomatal density of wheat and sour orange trees. *Journal of Experimental Botany*, **45**, 1665–1668.
- Farnsworth EJ, Ellison AM, Wong WK (1996) Elevated CO<sub>2</sub> alters anatomy, physiology, growth, and reproduction of red mangrove (*Rhizophora mangle* L.). *Oecologia*, **108**, 599–609.
- Farrar JF, Williams ML (1991) The effects of increased atmospheric carbon dioxide and temperature on carbon partitioning, source-sink relations and respiration. *Plant, Cell and Environment*, **14**, 819–830.
- Ferris R, Nijs I, Behaeghe T, Impens I (1996) Elevated CO<sub>2</sub> and temperature have different effects on leaf anatomy of perennial ryegrass in spring and summer. *Annals of Botany*, **78**, 489–497.
- Ferris R, Taylor G (1994) Increased root growth in elevated CO<sub>2</sub>: a biophysical analysis of root cell elongation. *Journal of Experimental Botany*, **280**, 1603–1612.
- Francis D (1992) The cell cycle in plant development. *New Phytologist*, **122**, 1–22.
- Garbutt K, Williams WE, Bazzaz FA (1990) Analysis of the differential response of five annuals to elevated CO<sub>2</sub> during growth. *Ecology*, **71**, 1185–1194.
- Gardner SDL, Bosac C, Taylor G (1995) Leaf growth of hybrid poplar following exposure to elevated CO<sub>2</sub>. *New Phytologist*, **131**, 81–90.
- Gaudillière JP, Mousseau M (1989) Short term effect of CO<sub>2</sub> enrichment on leaf development and gas exchange of young poplars (*Populus euramericana* cv I, 214). *Acta Oecologia: Oecologia Plantarum*, **10**, 95–105.
- Gloser J, Barták M (1994) Net photosynthesis, growth rate and biomass allocation in a rhizomatous grass *Calamagrostis epigejos* grown at elevated CO<sub>2</sub> concentration. *Photosynthetica*, **30**, 143–150.
- Goudriaan J, Zadoks JC (1995) Global climate change: Modeling the potential responses of agro-ecosystems with special reference to crop production. *Environmental Pollution*, **87**, 215–224.
- Gunderson CA, Wullschlegel SD (1994) Photosynthetic acclimation in trees to rising atmospheric CO<sub>2</sub>: a broader perspective. *Photosynthesis Research*, **39**, 369–388.
- Hirose T, Ackerly DD, Traw MB, Bazzaz FA (1996) Effects of CO<sub>2</sub> elevation on canopy development in the stands of two co-occurring annuals. *Oecologia*, **108**, 215–223.
- Ho LC (1977) Effects of CO<sub>2</sub> enrichment on the rates of photosynthesis and translocation of tomato leaves. *Annals of Applied Biology*, **87**, 191–200.
- Hunt R, Hand DW, Hannah MA, Neal AM (1991) Response to CO<sub>2</sub> enrichment in 27 herbaceous species. *Functional Ecology*, **5**, 410–421.
- Idso SB, Allen SG, Kimball BA (1990) Growth response of water lily to atmospheric CO<sub>2</sub> enrichment. *Aquatic Botany*, **37**, 87–92.

- Jacobs T (1997) Why do plant cells divide? *The Plant Cell*, **9**, 1021–1029.
- Jones HG (1985) Adaptive significance of leaf development and structural responses to environment. In: *Control of Leaf Growth* (eds Baker NR, Davies WJ, Ong CK), pp. 155–173. Cambridge University Press, Cambridge.
- Jongen M, Fay P, Jones MB (1996) Effects of elevated carbon dioxide and arbuscular mycorrhizal infection on *Trifolium repens*. *New Phytologist*, **132**, 413–423.
- Kendall AC, Turner JC, Thomas SM (1985) Effects of CO<sub>2</sub> enrichment at different irradiances on growth and yield of wheat. *Journal of Experimental Botany*, **36**, 252–260.
- Kende H, Zeevaert JAD (1997) The five 'classical' plant hormones. *The Plant Cell*, **9**, 1197–1210.
- Kerstetter RA, Hake S (1997) Shoot meristem formation in vegetative development. *The Plant Cell*, **9**, 1001–1010.
- Kinsman EA, Lewis C, Davies MS *et al.* (1996) Effects of temperature and elevated CO<sub>2</sub> on cell division in shoot meristems: differential response of two natural populations of *Dactylis glomerata* L. *Plant, Cell and Environment*, **19**, 775–780.
- Kinsman EA, Lewis C, Davies MS, Young JE, Francis D, Vilhar B, Ougham HJ (1997) Elevated CO<sub>2</sub> stimulates cells to divide in grass meristems: a differential effect in two natural populations of *Dactylis glomerata*. *Plant, Cell and Environment*, **20**, 1309–1316.
- Körner Ch (1991) Some overlooked plant characteristics as determinates of plant growth: a reconsideration. *Functional Ecology*, **5**, 162–173.
- Körner Ch, Bazzaz FA, Field CB (1996) The significance of biological variation, organism interactions, and life histories in CO<sub>2</sub> research. In: *Carbon Dioxide, Populations, and Communities* (eds Körner Ch, Bazzaz FA), pp. 443–455. Academic Press, San Diego, CA.
- Körner Ch, Palaez-Riedl S, Van Bel AJB (1995) CO<sub>2</sub> responsiveness of plants: a possible link to phloem loading. *Plant, Cell and Environment*, **18**, 595–600.
- Kouchi H, Sekine M, Hata S (1995) Distinct classes of mitotic cyclins are differentially expressed in the soybean shoot apex during the cell cycle. *The Plant Cell*, **7**, 1143–1155.
- Kozlowski TT, Pallardy SG (1997) *Growth Control in Woody Plants*. Academic Press, San Diego, CA.
- Küppers M (1985) Carbon relations and competition between woody species in a Central European hedgerow. III. Carbon and water balance on the leaf level. *Oecologia*, **65**, 94–100.
- Kutík J, Nátr L, Demmers-Derks HH, Lawlor DW (1995) Chloroplast ultrastructure of sugar beet (*Beta vulgaris* L.) cultivated in normal and elevated CO<sub>2</sub> concentrations with two contrasted nitrogen supplies. *Journal of Experimental Botany*, **46**, 1797–1802.
- Larigauderie A, Reynolds JF, Strain BR (1994) Root response to CO<sub>2</sub> enrichment and nitrogen supply in loblolly pine. *Plant and Soil*, **165**, 21–32.
- Leadley PW, Reynolds JF (1989) Effect of carbon dioxide enrichment on development of the first six mainstem leaves in soybean. *American Journal of Botany*, **76**, 1551–1556.
- Leadley PW, Reynolds JA, Thomas JF, Reynolds JF (1987) Effects of CO<sub>2</sub> enrichment on internal leaf surface area in soybeans. *Botanical Gazette*, **148**, 137–140.
- Lewis MC (1972) The physiological significance of variation in leaf structure. *Science Progress*, **60**, 25–51.
- Long SP (1991) Modification of the response of photosynthetic productivity to rising temperature by atmospheric CO<sub>2</sub> concentrations: has its importance been underestimated? *Plant, Cell and Environment*, **14**, 729–739.
- Long SP, Drake BG (1992) Photosynthetic CO<sub>2</sub> assimilation and rising atmospheric CO<sub>2</sub> concentrations. In: *Crop Photosynthesis: Spatial and Temporal Determinants* (eds Baker NR, Thomas H), pp. 69–107. Elsevier, New York.
- Madson E (1968) Effect of CO<sub>2</sub> concentration on the accumulation of starch and sugar in tomato leaves. *Physiologia Plantarum*, **21**, 168–175.
- McConnaughay KDM, Bernston GM, Bazzaz FA (1993) Plant responses to carbon dioxide. *Nature*, **361**, 24.
- Mendelsohn R, Rosenberg NJ (1994) Framework for integrated assessments of global warming impacts. *Climatic Change*, **28**, 15–44.
- Mo G, Nie D, Kirkham MB, He H, Ballou LK, Caldwell FW, Kanemasu ET (1992) Root and shoot weight in a tallgrass prairie under elevated carbon dioxide. *Environmental and Experimental Botany*, **32**, 193–201.
- Morison JIL, Gifford RM (1984a) Plant growth and water use with limited water supply in high CO<sub>2</sub> concentrations. I. Leaf area, water use and transpiration. *Australian Journal of Plant Physiology*, **11**, 361–374.
- Morison JIL, Gifford RM (1984b) Plant growth and water use with limited water supply in high CO<sub>2</sub> concentrations II. Plant dry weight, partitioning and water use efficiency. *Australian Journal of Plant Physiology*, **11**, 375–384.
- Mousseau M, Enoch HZ (1989) Carbon dioxide enrichment reduces shoot growth in sweet chestnut seedlings (*Castanea sativa* Mill.). *Plant, Cell and Environment*, **12**, 927–934.
- Murray DR (1995) Plant responses to carbon dioxide. *American Journal of Botany*, **82**, 690–697.
- Murray DR (1997) *Carbon Dioxide and Plant Responses*. John Wiley, Chichester, 275 pp.
- Murthy R, Dougherty PM (1997) Effect of carbon dioxide, fertilization and irrigation on loblolly pine branch morphology. *Trees*, **11**, 485–493.
- Newbery RM, Wolfenden J (1996) Effects of elevated CO<sub>2</sub> and nutrient supply on the seasonal growth and morphology of *Agrostis capillaris*. *New Phytologist*, **132**, 403–411.
- Nijs I, Impens I, Behaeghe T (1988) Effects of rising atmospheric carbon dioxide concentration on gas exchange and growth of perennial ryegrass. *Photosynthetica*, **22**, 44–50.
- Niklas KJ (1989) The effect of leaf-lobing on the interception of direct solar radiation. *Oecologia*, **80**, 59–64.
- Niklas KJ, Owens TG (1989) Physiological and morphological modifications of *Plantago major* (Plantaginaceae) in response to light conditions. *American Journal of Botany*, **76**, 370–382.
- Norby RJ (1996) Forest canopy productivity index. *Nature*, **381**, 6583.
- Norby RJ, O'Neill EG (1989) Growth dynamics and water use of seedlings of *Quercus alba* L. in CO<sub>2</sub>-enriched atmospheres. *New Phytologist*, **111**, 491–500.
- Norby RJ, O'Neill EG, Luxmoore RJ (1986) Effects of atmospheric CO<sub>2</sub> enrichment on the growth and mineral nutrition of *Quercus alba* seedlings in nutrient-poor soil. *Plant Physiology*, **82**, 83–89.
- O'leary JW, Knecht GN (1981) Elevated CO<sub>2</sub> concentration increases stomate numbers in *Phaseolus vulgaris* leaves. *Botanical Gazette*, **142**, 438–441.
- Oberbauer SF, Strain BR, Fetcher N (1985) Effect of CO<sub>2</sub>

- enrichment on seedling physiology and growth of two tropical tree species. *Physiological Planta*, **65**, 352–356.
- Pennanen A, Kemppe V, Lawlor D, Pehu E (1993) Effects of elevated CO<sub>2</sub> on photosynthesis, biomass production and chloroplast thylakoid structure of crop plants. *Current Topics in Plant Physiology*, **8**, 185–192.
- Pettersson R, McDonald AJS (1992) Effects of elevated carbon dioxide concentration on photosynthesis and growth of small birch plants (*Betula pendula* Roth.) at optimal nutrition. *Plant, Cell and Environment*, **15**, 911–919.
- Poethig RS (1997) Leaf morphogenesis in flowering plants. *The Plant Cell*, **9**, 1077–1087.
- Poorter H (1993) Interspecific variation in the growth response of plants to an elevated ambient CO<sub>2</sub> concentration. *Vegetatio*, **104/105**, 77–97.
- Poorter H, Van Berkel Y, Baxter R *et al.* (1997) The effect of elevated CO<sub>2</sub> on the chemical composition and construction costs of leaves of 27, C3 species. *Plant, Cell and Environment*, **20**, 472–482.
- Prior SA (1986) *Field studies of the water relations and growth responses of soybean [Glycine max (L.) Merr. 'Bragg'] grown under different water regimes in CO<sub>2</sub>-enriched atmospheres*. MS thesis, North Carolina State University, Raleigh, NC, 173 pp.
- Prior S, Runion GB, Mitchell RJ, Rogers HH, Amthor JS (1997) Effects of atmospheric CO<sub>2</sub> on longleaf pine: productivity and allocation as influenced by nitrogen and water. *Tree Physiology*, **17**, 397–405.
- Pritchard SG, Mosjidis C, Peterson CM, Runion GB, Rogers HH (1998) Anatomical and morphological alterations in longleaf pine needles resulting from growth in elevated CO<sub>2</sub>: interactions with soil resource availability. *International Journal of Plant Sciences*, **159**, 1002–1009.
- Pritchard SG, Peterson CM, Prior SA, Rogers HH (1997) Elevated atmospheric CO<sub>2</sub> differentially affects needle chloroplast ultrastructure and phloem anatomy in *Pinus palustris*: interactions with soil resource availability. *Plant, Cell and Environment*, **20**, 461–471.
- Pushnik JC, Demaree RS, Houpi JLJ, Flory WB, Bauer SM, Anderson PD (1995) The effect of elevated carbon dioxide on a Sierra-Nevadan dominant species: *Pinus ponderosa*. *Journal of Biogeography*, **22**, 249–254.
- Radoglou KM, Jarvis PG (1990a) Effects of CO<sub>2</sub> enrichment on four poplar clones. I. growth and leaf anatomy. *Annals of Botany*, **65**, 617–626.
- Radoglou KM, Jarvis PG (1990b) Effects of CO<sub>2</sub> enrichment on four poplar clones II. leaf surface properties. *Annals of Botany*, **65**, 627–632.
- Radoglou KM, Jarvis PG (1992) The effects of CO<sub>2</sub> enrichment and nutrient supply on growth morphology and anatomy of *Phaseolus vulgaris* L. seedlings. *Annals of Botany*, **70**, 245–256.
- Ranasinghe S, Taylor G (1996) Mechanism for increased leaf growth in elevated CO<sub>2</sub>. *Journal of Experimental Botany*, **47**, 349–358.
- Reekie EG (1996) The effect of elevated CO<sub>2</sub> on developmental processes and its implications for plant–plant interactions. In: *Carbon Dioxide, Populations, and Communities* (eds Körner Ch, Bazzaz FA ), pp. 333–346. Academic Press, San Diego, CA.
- Reekie EG, Bazzaz FA (1989) Competition and patterns of resource use among seedlings of five tropical trees grown at ambient and elevated CO<sub>2</sub>. *Oecologia*, **79**, 212–222.
- Renaudin JP, Colasanti J, Rime H, Yuan Z, Sundaresan V (1994) Cloning of four cyclins from maize indicates that higher plants have three structurally distinct groups of mitotic cyclins. *Proceedings of the National Academy of Science USA*, **91**, 7375–7379.
- Robertson EJ, Leech RM (1995) Significant changes in cell and chloroplast development in young wheat leaves (*Triticum aestivum* cv Hereward) grown in elevated CO<sub>2</sub>. *Plant Physiology*, **107**, 63–71.
- Robertson EJ, Williams M, Harwood JL, Lindsay J, Leaver JC, Leech RM (1995) Mitochondria increase three-fold and mitochondrial proteins and lipid change dramatically in postmeristematic cells in young wheat leaves grown in elevated CO<sub>2</sub>. *Plant Physiology*, **108**, 469–474.
- Rogers HH, Dahlman RC (1993) Crop responses to CO<sub>2</sub> enrichment. *Vegetatio*, **104/105**, 117–131.
- Rogers HH, Peterson CM, McCrimmon JN, Cure JD (1992) Response of plant roots to elevated atmospheric carbon dioxide. *Plant, Cell and Environment*, **15**, 749–752.
- Rogers HH, Prior SA, Runion GB, Mitchell RJ (1997a) Root to shoot ratio of crops as influenced by CO<sub>2</sub>. *Plant and Soil*, **187**, 229–248.
- Rogers HH, Runion GB, Krupa SV (1994) Plant responses to atmospheric CO<sub>2</sub> enrichment with emphasis on roots and the rhizosphere. *Environmental Pollution*, **83**, 155–189.
- Rogers HH, Runion GB, Krupa SV, Prior SA (1997b) Plant responses to atmospheric carbon dioxide enrichment: implications in root-soil-microbe interactions. In: *Advances in Carbon Dioxide Effects Research* (eds Allen LH, Kirkman MB, Olszyk DM, Whitman CE), pp. 1–34. ASA, Madison, WI.
- Rogers HH, Thomas JF, Bingham GE (1983) Response of agronomic and forest species to elevated atmospheric carbon dioxide. *Science*, **220**, 428–429.
- Ryle GJA, Powell CE (1992) The influence of elevated CO<sub>2</sub> and temperature on biomass production of continuously defoliated white clover. *Plant, Cell and Environment*, **15**, 593–599.
- Ryle GJA, Stanley J (1992) Effect of elevated CO<sub>2</sub> on stomatal size and distribution in perennial ryegrass. *Annals of Botany*, **69**, 563–565.
- Samuelson LJ, Seiler JR (1993) Interactive role of elevated CO<sub>2</sub> nutrient limitations, and water stress in the growth responses of red spruce seedlings. *Forest Science*, **39**, 348–358.
- Sasek TW, Strain BR (1989) Effects of carbon dioxide enrichment on the expansion and size of Kudzu (*Pueraria lobata*) leaves. *Weed Science*, **37**, 23–28.
- Sasek TW, Strain BR (1991) Effects of CO<sub>2</sub> enrichment on the growth and morphology of a native and an introduced honeysuckle vine. *American Journal of Botany*, **78**, 69–75.
- Sattler R, Rutishauser R (1997) The fundamental relevance of morphology and morphogenesis to plant research. *Annals of Botany*, **80**, 571–582.
- Saxe H, Ellsworth DS, Heath DS (1998) Tansley Review no. 98. Tree and forest functioning in an enriched CO<sub>2</sub> atmosphere. *New Phytologist*, **139**, 395–436.
- Schiefelbein JW, Masucci JD, Wang H (1997) Building a root: the control of patterning and morphogenesis during root development. *The Plant Cell*, **9**, 1089–1098.
- Sims DA, Seeman RJ, Luo Y (1998) Elevated CO<sub>2</sub> has independent effects on expansion rates and thickness of soybean leaves across light and nitrogen gradients. *Journal of Experimental Botany*, **49**, 583–591.
- Sionit N, Strain BR, Hellmers H, Riechers GH, Jaeger CH (1985)

- Long-term atmospheric CO<sub>2</sub> enrichment affects the growth and development of *Liquidambar styraciflua* and *Pinus taeda* seedlings. *Canadian Journal of Forest Research*, **15**, 468–471.
- Slafer GA, Rawson HM (1997) CO<sub>2</sub> effects on phasic development, leaf number and rate of leaf appearance in wheat. *Annals of Botany*, **79**, 75–81.
- Soni R, Carmichael JP, Shah ZH, Murray JAH (1995) A family of cyclin D homologues from plants differently controlled by growth regulators and containing the conserved retinoblastoma protein interaction motif. *Plant Cell*, **7**, 85–103.
- Spencer W, Bowes G (1986) Photosynthesis and growth of water hyacinth under CO<sub>2</sub> enrichment. *Plant Physiology*, **82**, 528–533.
- St. Omer L, Horvath SM (1984) Developmental changes in anatomy, morphology and biochemistry of *Layia platyglossa* exposed to elevated carbon dioxide. *American Journal of Botany*, **71**, 693–699.
- Stafstrom JP (1995) Developmental potential of shoot buds. In: *Plant Stems: Physiological and Functional Morphology* (ed. Gartner BL.), pp. 257–279 Academic Press, San Diego, CA.
- Stitt M (1991) Rising CO<sub>2</sub> levels and their potential significance for carbon flow in photosynthetic cells. *Plant, Cell and Environment*, **14**, 741–762.
- Stulen I, den Hertog J (1993) Root growth and functioning under atmospheric CO<sub>2</sub> enrichment. *Vegetatio*, **104/105**, 99–115.
- Taylor CB (1997) Plant vegetative development: from seed and embryo to shoot and root. *The Plant Cell*, **9**, 981–988.
- Taylor G, Ranasinghe S, Bosac C, Gardner SDL, Ferris R (1994) Elevated CO<sub>2</sub> and plant growth: cellular mechanisms and responses of whole plants. *Journal of Experimental Botany*, **45**, 1761–1774.
- Teskey RO, Bongarten BC, Cregg BM, Dougherty PM, Hennessey TC (1987) Physiology and genetics of tree growth response to moisture and temperature stress: an examination of the characteristics of loblolly pine (*Pinus taeda* L.). *Tree Physiology*, **3**, 41–61.
- Teugels H, Nijs I, Van Hecke P, Impens I (1995) Competition in a global change environment: the importance of different plant traits for competitive success. *Journal of Biogeography*, **22**, 297–305.
- Thomas SC, Bazzaz FA (1996) Elevated CO<sub>2</sub> and leaf shape: are dandelions getting toothier? *American Journal of Botany*, **83**, 106–111.
- Thomas JF, Harvey CH (1983) Leaf anatomy of four species grown under continuous CO<sub>2</sub> enrichment. *Botanical Gazette*, **144**, 303–309.
- Ticha I (1982) Photosynthetic characteristics during ontogenesis of leaves. 7. Stomata density and sizes. *Photosynthetica*, **16**, 375–471.
- Tolley LC, Strain BR (1984) Effects of CO<sub>2</sub> enrichment on growth of *Liquidambar styraciflua* and *Pinus taeda* seedlings under different irradiance levels. *Canadian Journal of Forestry Research*, **14**, 343–350.
- Tremmel DC, Bazzaz FA (1993) How neighbor canopy architecture affects target plant performance. *Ecology*, **74**, 2114–2124.
- Tyree MT, Alexander JD (1993) Plant water relations and the effects of elevated CO<sub>2</sub>: a review and suggestions for further research. *Vegetatio*, **104/105**, 47–62.
- Van Oosten JJ, Wilkens D, Besford RT (1994) Regulation of the expression of photosynthetic nuclear genes by CO<sub>2</sub> is mimicked by regulation by carbohydrates: a mechanism for the acclimation of photosynthesis to high CO<sub>2</sub>? *Plant, Cell and Environment*, **17**, 913–923.
- Vu JCV, Allen LH, Bowes G (1989) Leaf ultrastructure, carbohydrates and protein of soybeans grown under CO<sub>2</sub> enrichment. *Environmental and Experimental Botany*, **29**, 141–147.
- Wagner F, Below R, De Klerk P, Dilcher DL, Joosten H, Kureschner WM, Visscher H. (1996) A natural experiment on plant acclimation: lifetime stomatal frequency response of an individual tree to annual atmospheric CO<sub>2</sub> increase. *Proceedings of the National Academy of Sciences U.S.A.*, **93**, 11,705–11,708.
- Weber JA, Grulke NE (1995) Response of stem growth and function to air pollution. In: *Plant Stems: Physiological and Functional Morphology* (ed. Gartner BL.), pp. 343–363 Academic Press, San Diego, CA.
- Wittwer SH (1995) Food, Climate, and Carbon Dioxide. *The Global Environment and World Food Production*. CRC Press, Boca Raton, FL.
- Wong SC (1979) Elevated atmospheric partial pressure of CO<sub>2</sub> and plant growth I. Interactions of nitrogen nutrition and photosynthetic capacity in C<sub>3</sub> and C<sub>4</sub> plants. *Oecologia*, **44**, 68–74.
- Woodward FI, Bazzaz FA (1988) The responses of stomatal density to CO<sub>2</sub> partial pressure. *Journal of Experimental Botany*, **39**, 1771–1781.
- Wulff RD, Strain BR (1981) Effects of CO<sub>2</sub> enrichment on growth and photosynthesis in *Desmodium paniculatum*. *Canadian Journal of Botany*, **60**, 1084–1091.
- Yegappan TM, Paton DM, Gates CT (1982) Water stress in sunflower (*Helianthus annuus* L.) 2. Effects on leaf cells and leaf area. *Annals of Botany*, **49**, 63–68.
- Yelle S, Beeson RC, Trudel MJ, Gosselin A (1989) Acclimation of two tomato species to high atmospheric CO<sub>2</sub>. I. Starch and sugar concentrations. *Plant Physiology*, **90**, 1465–1472.
- Zimmermann MH (1983) Hydraulic architecture of some diffuse porous trees. *Canadian Journal of Botany*, **56**, 2286–2295.